

UNGUARDED RECURSION ON COINDUCTIVE RESUMPTIONS*

SERGEY GONCHAROV, CHRISTOPH RAUCH, LUTZ SCHRÖDER, AND JULIAN JAKOB

Department of Computer Science, Friedrich-Alexander-Universität Erlangen-Nürnberg
e-mail address: {sergey.goncharov,christoph.rauch,lutz.schroeder,julian.jakob}@fau.de

ABSTRACT. We study a model of side-effecting processes obtained by starting from a monad modelling base effects and adjoining free operations using a cofree coalgebra construction; one thus arrives at what one may think of as types of non-wellfounded side-effecting trees, generalizing the infinite resumption monad. Correspondingly, the arising monad transformer has been termed the coinductive generalized resumption transformer. Monads of this kind have received some attention in the recent literature; in particular, it has been shown that they admit guarded iteration. Here, we show that they also admit unguarded iteration, i.e. form complete Elgot monads, provided that the underlying base effect supports unguarded iteration. Moreover, we provide a universal characterization of the coinductive resumption monad transformer in terms of coproducts of complete Elgot monads.

1. INTRODUCTION

Following seminal work by Moggi [24], monads are widely used to represent computational effects in program semantics, and in fact in actual programming languages [35]. Their main attraction lies in the fact that they provide an interface to a generic notion of side-effect at the right level of abstraction: they subsume a wide variety of side-effects such as state, non-determinism, random, and I/O, and at the same time retain enough internal structure to support a substantial amount of generic meta-theory and programming, the latter witnessed, for example, by the monad class implemented in the Haskell basic libraries [25].

In the current work, we study a particular construction on monads motivated partly by the goal of modelling generic side-effects in the semantics of reactive processes. Specifically, given a base monad T and a strong functor Σ , we have final coalgebras

$$T_{\Sigma}X = \nu\gamma. T(X + \Sigma\gamma)$$

for each object X , assuming enough structure on T , Σ , and the base category. Inhabitants of $T_{\Sigma}X$ are understood as (possibly) nonterminating processes that proceed in steps, where each step produces side-effects specified by T (e.g. writing to shared global memory, non-determinism) and performs communication actions specified by Σ . E.g. in the simplest

1998 ACM Subject Classification: F.3.2 [Logics and Meanings of Programs]: Semantics of Programming Languages — algebraic approaches to semantics, denotational semantics; F.3.3 [Logics and Meanings of Programs]: Studies of Program Constructs — program and recursion schemes; D.3.3 [Programming languages]: Language Constructs and Features — recursion; General Terms: Theory.

Key words and phrases: Recursion, coalgebra, coinduction, complete Elgot monad, resumptions.

* Extended and revised version of [16].

case, Σ is of the form $a \times (-)^b$, which may be understood as reading inputs of type b and writing outputs of type a .

The construction of $T_\Sigma X$ from T is an infinite version of the generalized resumption transformer introduced by Cienciarelli and Moggi [13]. It has been termed the *coinductive generalized resumption* transformer by Piróg and Gibbons [26, 27], who show that on the Kleisli category of T , T_Σ is the free completely iterative monad generated by $T(a \times _^b)$.

The result that T_Σ is a completely iterative monad brings us to the contribution of the current paper. Recall that complete iterativity of T_Σ means that for every morphism

$$e : X \rightarrow T_\Sigma(Y + X),$$

read as an equation defining the inhabitants of X , thought of as variables, as terms over the defined variables (from X) and parameters from Y , has a unique *solution*

$$e^\dagger : X \rightarrow T_\Sigma Y$$

in the evident sense, *provided* that e is *guarded*. The latter concept is defined in terms of additional structure of T_Σ as an *idealized monad*, which essentially allows distinguishing terms beginning with an operation from mere variables. Guardedness of e then means that recursive calls can happen only under a free operation. Similar results on guarded recursion abound in the literature; for example, the fact that T_Σ admits guarded recursive definitions can also be deduced from more general results by Uustalu on parametrized monads [34].

The central result of the current paper is to remove the guardedness restriction in the above setup. That is, we show that a solution $e^\dagger : X \rightarrow T_\Sigma Y$ exists for *every* morphism $e : X \rightarrow T_\Sigma(X + Y)$. Of course, the solution is then no longer unique (for example, we admit definitions of the form $x = x$); moreover, we clearly need to make additional assumptions about T . Our result states, more precisely, that T_Σ allows for a principled *choice* of solutions $e^\dagger$ satisfying standard equational laws for recursion [32], thus making T_Σ into a *complete Elgot monad* [3]¹. The assumption on T that we need to enable this result is that T itself is an Elgot monad (e.g. partiality, nondeterminism, or combinations of these with state), i.e. we show that *the class of Elgot monads is stable under the coinductive generalized resumption transformer*. We show moreover that the structure of T_Σ as an Elgot monad is uniquely determined as extending that of T .

The motivation for these results is, well, to free non-wellfounded recursive definitions from the standard guardedness constraint. Note for example that in [26], it was necessary to assume guards in all loop iterations when interpreting a while-language with actions originally proposed by Rutten [31] over a completely iterative monad. Contrastingly, given that T_Σ is a (complete) Elgot monad, one can now just write unrestricted while loops. We elaborate this example in Section 7, and recall a standard example of unguarded recursion in process algebra in Section 8.

An earlier version of this work has appeared as [16]; the present version not only has full proofs, but also works in a generalized setup with an arbitrary strong functor Σ (admitting the requisite final coalgebras) instead of just functors of the form $a \times (-)^b$.

¹We modify the original definition of Elgot monad, which requires the object X of variables to be a finitely presentable object in an lfp category, by admitting unrestricted objects of variables. This change is owed mostly to the fact that we do not assume the base category to be lfp, and in our own estimate appears to be technically inessential, although we have not checked details for the obvious variants of our results that arise by replacing complete Elgot monads with Elgot monads.

2. PRELIMINARIES

According to Moggi [23], a notion of computation can be formalized as a strong monad $\mathbb{T}$ over a Cartesian category (i.e. a category with finite products). In order to support the constructions occurring in the main object of study, we work in a *distributive category* $\mathbf{C}$, i.e. a category with finite products and coproducts (including a final and an initial object) such that the natural transformation

$$X \times Y + X \times Z \xrightarrow{[\text{id} \times \text{inl}, \text{id} \times \text{inr}]} X \times (Y + Z)$$

is an isomorphism [14], whose inverse we denote $\text{dist}_{X,Y,Z}$. Here we denote injections into binary coproducts by $\text{inl} : X \rightarrow X + Y$, $\text{inr} : Y \rightarrow X + Y$. The projections from binary products are denoted $\text{fst} : X \times Y \rightarrow X$, $\text{snd} : X \times Y \rightarrow Y$; pairing is denoted by $\langle _, _ \rangle$, and copairing of $f : X \rightarrow Z$, $g : Y \rightarrow Z$ by $[f, g] : X + Y \rightarrow Z$. Unique morphisms $A \rightarrow 1$ into the terminal object are written $!_X$, or just $!$. We write $|\mathbf{C}|$ for the class of objects of $\mathbf{C}$. Distributivity essentially allows using context variables in case expressions, i.e. in copairing. We omit indices on natural transformations where this is unlikely to cause confusion.

A *strong* functor on $\mathbf{C}$ is a functor $F : \mathbf{C} \rightarrow \mathbf{C}$ equipped with a natural transformation

$$\rho_{X,Y} : X \times FY \rightarrow F(X \times Y)$$

called *strength*, subject to the equations

$$\text{snd} = (F \text{snd}) \rho \tag{STR_1}$$

$$(F \text{assoc}) \rho = \tau(\text{id} \times \rho) \text{assoc} \tag{STR_2}$$

where $\text{assoc} : (X \times Y) \times Z \rightarrow X \times (Y \times Z)$ is the associativity isomorphism of products, explicitly, $\text{assoc} = \langle \text{fst} \text{fst}, \langle \text{snd} \text{fst}, \text{snd} \rangle \rangle$. A natural transformation $\alpha : F \rightarrow G$ between strong functors F, G (with the strength denoted ρ in both cases) is *strong* if it commutes with strength:

$$\begin{array}{ccc} X \times FY & \xrightarrow{\text{id}_X \times \alpha_Y} & X \times GY \\ \rho_{X,Y} \downarrow & & \downarrow \rho_{X,Y} \\ F(X \times Y) & \xrightarrow{\alpha_{X \times Y}} & G(X \times Y) \end{array}$$

Recall that a monad $\mathbb{T}$ over $\mathbf{C}$ can be given by a *Kleisli triple* $(T, \eta, _*)$ where T is an endomap of $|\mathbf{C}|$ (in the following, we always denote monads and their functor parts by the same letter, with the former in blackboard bold), the *unit* η is a family of morphisms $\eta_X : X \rightarrow TX$, and the *Kleisli lifting* $_*$ maps $f : X \rightarrow TY$ to $f^* : TX \rightarrow TY$, subject to the equations

$$\eta^* = \text{id} \qquad f^* \eta = f \qquad (f^* g)^* = f^* g^*.$$

This is equivalent to the presentation in terms of an endofunctor T with natural transformations unit and multiplication.

A *strong monad* is a monad equipped with a strength τ satisfying the following additional coherence conditions [23] (with modifications reflecting the switch from monad multiplication to Kleisli lifting):

$$\tau(\text{id} \times \eta) = \eta \tag{STR_3}$$

$$(\tau(\text{id} \times f))^* \tau = \tau(\text{id} \times f^*) \tag{STR_4.}$$

The typing of the law (STR₄) capturing compatibility of the strength with Kleisli lifting is shown in the diagram

$$\begin{array}{ccc}
 X \times TY & \xrightarrow{\text{id} \times f^*} & X \times TZ \\
 \tau \downarrow & & \downarrow \tau \\
 T(X \times Y) & \xrightarrow{(\tau(\text{id} \times f))^*} & T(X \times Z)
 \end{array}$$

$$X \times Y \xrightarrow{\text{id} \times f} X \times TZ \xrightarrow{\tau} T(X \times Z)$$

(For distinction, we denote strengths of monads by τ and strengths of functors by ρ throughout.) Strength enables interpreting programs over more than one variable, and allows for internalization of the Kleisli lifting, thus legitimating expressions like $\lambda x. (f(x))^* : X \rightarrow (TY \rightarrow TZ)$ for $f : X \rightarrow (Y \rightarrow TZ)$, which encodes $\text{curry}(\text{uncurry}(f)^* \tau)$. Strength is equivalent to the monad being enriched over $\mathbf{C}$ [20]; in particular, every monad on **Set** is strong. Henceforth we shall use the term ‘monad’ to mean ‘strong monad’ unless explicitly stated otherwise. We emphasize however that all our results remain valid under the removal of all strength assumptions and claims (that is, replacing the terms strong monad, strong functor, and strong natural transformation with monad, functor, and natural transformation, respectively, throughout).

The standard intuition for a monad $\mathbb{T}$ is to think of TX as the set of terms in some algebraic theory, with variables taken from X . In this view, the unit converts variables into terms, and a Kleisli lifting f^* applies a substitution $f : X \rightarrow TY$ to terms over X . In our setting, the ‘terms’ featuring here are often infinite; nevertheless, we sometimes call them *algebraic terms* for emphasis.

The *Kleisli category* $\mathbf{C}_{\mathbb{T}}$ of a monad $\mathbb{T}$ has the same objects as $\mathbf{C}$, and $\mathbf{C}$ -morphisms $X \rightarrow TY$ as morphisms $X \rightarrow Y$. The identity on X in $\mathbf{C}_{\mathbb{T}}$ is η_X ; and the *Kleisli composite* of $f : X \rightarrow TY$ and $g : Y \rightarrow TZ$ is $g^* f$. A monad $\mathbb{T}$ has *rank* κ if it preserves κ -filtered colimits. On **Set**, this condition intuitively means that T is determined by its values on sets of cardinality less than κ .

Example 2.1. The main motivating example for Σ is obtained as follows. Assume that $\mathbf{C}$ has *exponentials* of the form X^b (for b ranging over a subset of $|\mathbf{C}|$), i.e. objects adjoint to Cartesian products $X \times b$, which means that for any X and Y , there is an isomorphism

$$\text{curry}_{X,Y} : \text{Hom}_{\mathbf{C}}(X \times b, Y) \cong \text{Hom}_{\mathbf{C}}(X, Y^b),$$

natural in X and Y . We write $\text{uncurry}_{X,Y}$ for the inverse map $\text{curry}_{X,Y}^{-1}$. The *evaluation morphism* $\text{ev}_X : X^b \times b \rightarrow X$ (natural in X) is obtained as $\text{uncurry}_{X^b, X}(\text{id}_{X^b})$.

It is easy to see that the functors $X \mapsto a \times X$ and $X \mapsto X^b$ are strong and that composites and coproducts of strong functors (as plain functors) are again strong functors. Hence, the functor

$$\Sigma X = \sum_i a_i \times X^{b_i}$$

is strong. Intuitively (and formally correctly on **Set**), ΣX can be seen as the set of flat terms over variables from X in the signature Σ , i.e. the elements of ΣX are of the form $f_i(c; x_1, \dots, x_{n_i})$ where c is a *parameter* from a_i and $x_1, \dots, x_{n_i}$ are elements of X . The

computational meaning of exponents in X^b is thus to capture a notion of arity of algebraic operations generating effects, e.g. $b = 2$ would correspond to binary operations such as nondeterministic choice. An example of an operation taking a parameter would be the operation of writing a value val to position ind of an array, $update(\langle ind, val \rangle; x)$ (see [29] for details).

A more general setup involves categories enriched over a symmetric monoidal closed category $\mathbf{V}$ whose objects are then treated as arities (and *coarities*, i.e. objects used for indexing families of operations) [19, 18]. One then replaces products with *tensors* and exponentials with *cotensors*.

Another example are functors on the category **Nom** of nominal sets and equivariant maps built using constant functors, identity, coproducts, finite products, and the so-called *abstraction functor* $[\mathbb{A}](-)$, where $\mathbb{A}$ is a set of *names* and $[\mathbb{A}]X$ consists of pairs $(a, x) \in \mathbb{A} \times X$ modulo a natural notion of α -equivalence [28]. Such functors represent so-called *binding signatures*, whose operations may bind names, such as λ -abstraction or π -calculus-style fresh name binders ν ; terms are then taken modulo α -equivalence. E.g. the syntax of the λ -calculus is generated as the initial algebra of the functor $LX = \mathbb{A} + X \times X + [\mathbb{A}]X$ (see [15]).

3. COMPLETE ELGOT MONADS

As indicated in the introduction, we will be interested in recursive definitions over a monad $\mathbb{T}$; abstractly, these are morphisms

$$f : X \rightarrow T(Y + X)$$

thought of as associating to each variable $x : X$ a definition $f(x)$ in the shape of an algebraic term from $T(Y + X)$, which thus employs parameters from Y as well as the defined variables from X . The latter amount to recursive calls of the definition. This notion is agnostic to what happens in the case of non-terminating recursion. For example, T might identify all non-terminating sequences of recursive calls into a single value $\perp$ signifying non-termination; at the other extreme, T might be a type of infinite trees that just records the tree of recursive calls explicitly.

To a recursive definition f as above, we wish to associate a solution

$$f^\dagger : X \rightarrow TY,$$

which amounts to a non-recursive definition of the elements of X as terms over Y only. As we do not assume any form of guardedness, this solution will in general fail to be unique. We thus require a coherent selection of solutions $f^\dagger$ for all equations f , where by coherent we mean that the selection satisfies a collection of well-established (quasi-)equational properties. Formally:

Definition 3.1. (Complete Elgot monads) A *complete Elgot monad* is a monad $\mathbb{T}$ equipped with an operator $_^\dagger$, called *iteration*, that assigns to each morphism $f : X \rightarrow T(Y + X)$ a morphism $f^\dagger : X \rightarrow TY$ such that the following conditions hold:

- *unfolding*: $[\eta, f^\dagger]^* f = f^\dagger$;
- *naturality*: $g^* f^\dagger = ([T \text{ inl } g, \eta \text{ inr}]^* f)^\dagger$ for $g : Y \rightarrow TZ$;
- *dinaturality*: $([\eta \text{ inl}, h]^* g)^\dagger = [\eta, ([\eta \text{ inl}, g]^* h)^\dagger]^* g$ for $g : X \rightarrow T(Y + Z)$ and $h : Z \rightarrow T(Y + X)$;

- *codiagonal*: $(T[\text{id}, \text{inr}] g)^\dagger = (g^\dagger)^\dagger$ for $g : X \rightarrow T((Y + X) + X)$;
- *uniformity*: $f h = T(\text{id} + h) g$ implies $f^\dagger h = g^\dagger$ for $g : Z \rightarrow T(Y + Z)$ and $h : Z \rightarrow X$.

Additionally, iteration must be *compatible with strength* in the sense that

$$\tau(\text{id} \times f^\dagger) = (T \text{ dist } \tau(\text{id} \times f))^\dagger$$

for $f : X \rightarrow T(Y + X)$.

Remark 3.2. The above definition is inspired by the axioms of parametrized uniform iterativity [32], which go back to Bloom and Ésik [11]. Adámek et al. [3] define *Elgot monads* by means of a slightly different system of axioms: the codiagonal and dinaturality axioms are replaced with the *Bekić identity*. Both axiomatizations are however equivalent, which is essentially a result about iteration theories [11, Section 6.8]; we record a self-contained proof of this equivalence in Proposition 3.3 below. Moreover, the iteration operator in [3] is defined only for $f : X \rightarrow T(Y + X)$ with finitely presentable X , under the assumption that $\mathbf{C}$ is *locally finitely presentable*; hence our use of the term ‘complete Elgot monad’ instead of ‘Elgot monad’. We have the impression that this difference is not technically essential but have not checked details for the finitary variant of our results.

Proposition 3.3 (Bekić identity). *A complete Elgot monad $\mathbb{T}$ is equivalently a monad satisfying unfolding, naturality, uniformity (as in Definition 3.1), and the Bekić identity*

$$(T\alpha[f, g])^\dagger = [\eta, h^\dagger]^\star [\eta \text{ inr}, g^\dagger] \quad (\text{Bekić})$$

where $g : X \rightarrow T((Z + Y) + X)$, $f : Y \rightarrow T((Z + Y) + X)$, $h = [\eta, g^\dagger]^\star f : Y \rightarrow T(Z + Y)$, with $\alpha : (A + B) + C \rightarrow A + (B + C)$ being the obvious coproduct associativity morphism.

Proof. Let us show that complete Elgot monads validate the Bekić identity. Let

$$u = T((\text{id} + \text{inl}) + \text{inr})[f, g] : Y + X \rightarrow T((Z + (Y + X)) + (Y + X)).$$

By *codiagonal*,

$$(T[\text{id}, \text{inr}] u)^\dagger = (u^\dagger)^\dagger. \quad (3.1)$$

Now the left-hand side of (3.1) simplifies to

$$\begin{aligned} & (T[\text{id}, \text{inr}] T((\text{id} + \text{inl}) + \text{inr})[f, g])^\dagger \\ &= (T[\text{id} + \text{inl}, \text{inr inr}][f, g])^\dagger \\ &= (T\alpha[f, g])^\dagger, \end{aligned}$$

i.e. to the left-hand side of the Bekić identity. Now observe that, by *uniformity* and *naturality*,

$$u^\dagger \text{ inr} = (T(\text{id} + \text{inl}) + \text{id}) g^\dagger = T(\text{id} + \text{inl}) g^\dagger. \quad (3.2)$$

Therefore, the right-hand side of (3.1) can be rewritten in the form

$$\begin{aligned}
 (u^\dagger)^\dagger &= ([\eta, u^\dagger]^\star u)^\dagger && \text{(unfolding)} \\
 &= ([\eta(\text{id} + \text{inl}), u^\dagger \text{inr}][f, g])^\dagger \\
 &= ([T(\text{id} + \text{inl}) \eta, T(\text{id} + \text{inl}) g^\dagger]^\star [f, g])^\dagger && (3.2) \\
 &= (T(\text{id} + \text{inl})[\eta, g^\dagger]^\star [f, g])^\dagger \\
 &= (T(\text{id} + \text{inl})[[\eta, g^\dagger]^\star f, g^\dagger])^\dagger && \text{(unfolding)} \\
 &= ([\eta \text{inl}, \eta \text{inr inl}]^\star [[\eta, g^\dagger]^\star f, g^\dagger])^\dagger \\
 &= [\eta, ([\eta \text{inl}, [[\eta, g^\dagger]^\star f, g^\dagger]]^\star \eta \text{inr inl})^\dagger]^\star \\
 &\quad [[\eta, g^\dagger]^\star f, g^\dagger] && \text{(dinaturality)} \\
 &= [\eta, ([\eta, g^\dagger]^\star f)^\dagger]^\star [[\eta, g^\dagger]^\star f, g^\dagger] \\
 &= [[[\eta, g^\dagger]^\star f]^\dagger, [\eta, ([\eta, g^\dagger]^\star f)^\dagger]^\star g^\dagger] && \text{(unfolding)} \\
 &= [h^\dagger, [\eta, h^\dagger]^\star g^\dagger] \\
 &= [\eta, h^\dagger]^\star [\eta \text{inr}, g^\dagger],
 \end{aligned}$$

i.e. equals the right-hand side of the Bekić identity.

For the opposite direction, we need to show that the Bekić identity implies *dinaturality* and *codiagonal*. For the latter, let $k : X \rightarrow T((Y + X) + X)$. By the Bekić identity,

$$\begin{aligned}
 (T\alpha[k, k])^\dagger &= [\eta, ([\eta, k^\dagger]^\star k)^\dagger]^\star [\eta \text{inr}, k^\dagger] \\
 &= [\eta, (k^\dagger)^\dagger]^\star [\eta \text{inr}, k^\dagger].
 \end{aligned}$$

Thus, $(T\alpha[k, k])^\dagger \text{inl} = (T\alpha[k, k])^\dagger \text{inr} = (k^\dagger)^\dagger$. On the other hand, by *uniformity*,

$$(T\alpha[k, k])^\dagger = (T[\text{id}, \text{inr}] k)^\dagger [\text{id}, \text{id}]$$

and therefore

$$(k^\dagger)^\dagger = (T\alpha[k, k])^\dagger \text{inr} = (T[\text{id}, \text{inr}] k)^\dagger$$

as required. To prove *dinaturality*, we define the term

$$w = ([T(\text{id} + \text{inr}) h, T(\text{id} + \text{inl}) g])^\dagger$$

for $g : X \rightarrow T(Y + Z)$, $h : Z \rightarrow T(Y + X)$. By *uniformity*,

$$w[\text{inr}, \text{inl}] = ([T(\text{id} + \text{inr}) g, T(\text{id} + \text{inl}) h])^\dagger.$$

The Bekić identity then gives us

$$\begin{aligned}
 w &= ([T(\text{id} + \text{inr}) h, T(\text{id} + \text{inl}) g])^\dagger \\
 &= (T\alpha[T(\text{inl} + \text{id}) h, T \text{inl} g])^\dagger \\
 &= [\eta, ([\eta, (T \text{inl} g)^\dagger]^\star T(\text{inl} + \text{id}) h)^\dagger]^\star [\eta \text{inr}, (T \text{inl} g)^\dagger] \\
 &= [\eta, ([\eta \text{inl}, g]^\star h)^\dagger]^\star [\eta \text{inr}, g]
 \end{aligned}$$

as well as

$$w[\text{inr}, \text{inl}] = [\eta, ([\eta \text{inl}, h]^\star g)^\dagger]^\star [\eta \text{inr}, h].$$

Therefore,

$$([\eta \text{ inl}, h]^\star g)^\dagger = w[\text{inr}, \text{inl}] \text{ inl} = w \text{ inr} = [\eta, ([\eta \text{ inl}, g]^\star h)^\dagger]^\star g. \quad \square$$

Given a complete Elgot monad $\mathbb{T}$, we can parametrize the iteration operator $_^\dagger$ with an additional argument to be carried over the recursion loop, i.e. we derive an operator $_^\ddagger$ sending $f : Z \times X \rightarrow T(Y + X)$ to $f^\ddagger : Z \times X \rightarrow TY$ by

$$f^\ddagger = (T(\text{snd} + \text{id})(T \text{ dist}) \tau_{Z, Y+X} \langle \text{fst}, f \rangle)^\dagger. \quad (3.3)$$

We call the derived operator $_^\ddagger$ *strong iteration*.

The key examples of complete Elgot monads are, on the one hand, so-called ω -continuous monads (Definition 3.4), and, on the other hand, extensions of complete Elgot monads, e.g. of ω -continuous monads, with free operations. The latter arise by application of the coinductive generalized resumption transformer as introduced in Section 4. We proceed to discuss ω -continuous monads, which are defined as having a suitable order-enrichment of their Kleisli category.

Definition 3.4. (ω -continuous monad) An ω -continuous monad consists of a monad $\mathbb{T}$ and an enrichment of the Kleisli category $\mathbf{C}_\mathbb{T}$ of $\mathbb{T}$ over the category $\omega\mathbf{Cppo}$ of ω -complete partial orders with bottom and (nonstrict) continuous maps, satisfying the following conditions:

- strength is ω -continuous: $\tau(\text{id} \times \bigsqcup_i f_i) = \bigsqcup_i (\tau(\text{id} \times f_i))$;
- copairing in $\mathbf{C}_\mathbb{T}$ is ω -continuous in both arguments: $[\bigsqcup_i f_i, \bigsqcup_i g_i] = \bigsqcup_i [f_i, g_i]$;
- bottom elements are preserved by strength and by postcomposition in $\mathbf{C}_\mathbb{T}$: $\tau(\text{id} \times \perp) = \perp$, $f^\star \perp = \perp$.

Example 3.5. Many of the standard computational monads on **Set** [23] are ω -continuous, including nontermination ($TX = X + 1$), nondeterminism ($TX = \mathcal{P}(X)$), and the non-deterministic state monad ($TX = \mathcal{P}(X \times S)^S$ for a set S of states). On $\omega\mathbf{Cppo}$, lifting ($TX = X_\perp$) and the various power domain monads are ω -continuous.

Remark 3.6. As observed by Kock [20], monad strength is equivalent to enrichment over the base category. One consequence of this fundamental fact is that if $\mathbf{C}$ is enriched over the category $\omega\mathbf{Cpo}$ of bottomless ω -complete partial orders and ω -continuous maps (i.e. $\mathbf{C}$ is an **O**-category in the sense of Wand [36] and of Smyth and Plotkin [33]), with the bicartesian closed structure enriched in the obvious sense, then $\mathbf{C}_\mathbb{T}$ is also enriched over $\omega\mathbf{Cpo}$, since T , being a strong functor, is an $\omega\mathbf{Cpo}$ -functor (aka *locally continuous* functor [33]). Then $\mathbb{T}$ is ω -continuous in the sense of Definition 3.4 iff each $\text{Hom}(X, TY)$ has a bottom element preserved by strength and postcomposition in $\mathbf{C}_\mathbb{T}$. This allows for incorporating numerous domain-theoretic examples by taking $\mathbf{C}$ to be a suitable category of predomains, and $\mathbb{T}$, in the simplest case, the *lifting monad* $TX = X_\perp$ (from which one builds more complex examples by the construction explored next).

If $\mathbb{T}$ is an ω -continuous monad, then the endomap

$$h \mapsto [\eta, h]^\star f \quad (3.4)$$

on the hom-set $\text{Hom}_\mathbf{C}(A, TB)$ is continuous because copairing and Kleisli composition in T are continuous, and hence has a least fixpoint by Kleene's fixpoint theorem. We can define an iteration operator by taking $f^\dagger$ to be this fixpoint; in other words, $f^\dagger$ is defined to be the smallest solution of the unfolding equation as per Definition 3.1. This yields

Theorem 3.7. *On every ω -continuous monad, defining iteration by taking least fixpoints determines a complete Elgot monad structure.*

This result is to be expected in the light of analogous facts known for Bloom and Ésik's ω -continuous theories [11, Theorem 8.2.15, Exercise 8.2.17].

Proof. Let $\mathbb{T}$ be an ω -continuous monad, and let $f^\dagger$ be the least fixpoint of (3.4).

Let us verify the axioms of Elgot monads one by one. To that end we employ the following *uniformity rule* for least fixpoints of continuous functionals [32]:

$$\frac{UF = GU \quad U(\perp) = \perp}{U(\mu F) = \mu G} \quad (3.5)$$

- *Unfolding.* This holds by definition.
- *Naturality.* In (3.5) take $F(u) = [\eta, u]^* f$, $G(u) = [\eta, u]^* [(T \text{ inl})g, \eta \text{ inr}]^* f$ and $U(u) = g^* u$. By definition, $U(\perp) = \perp$, $\mu F = f^\dagger$, $\mu G = ([\eta, f^\dagger]^* f)^\dagger$. Then we have

$$U(F(u)) = g^* [\eta, u]^* f = [\eta, g^* u]^* [(T \text{ inl})g, \eta \text{ inr}]^* f = G(U(u)).$$

Therefore, by (3.5), $g^* f^\dagger = U(\mu F) = \mu G = ([\eta, f^\dagger]^* f)^\dagger$.

- *Dinaturality.* Let us denote $s = [\eta \text{ inl}, h]^* g$ and $t = [\eta \text{ inl}, g]^* h$. The identity in question is then $s^\dagger = [\eta, t^\dagger]^* g$. Observe that

$$\begin{aligned} [\eta, t^\dagger]^* g &= [\eta, [\eta, t^\dagger]^* [\eta \text{ inl}, g]^* h]^* g && \text{(unfolding)} \\ &= [\eta, [\eta, [\eta, t^\dagger]^* g]^* h]^* g \\ &= [\eta, [\eta, t^\dagger]^* g]^* [\eta \text{ inl}, h]^* g, \end{aligned}$$

i.e. $[\eta, t^\dagger]^* g$ satisfies the unfolding identity for $s^\dagger$, therefore $s^\dagger \sqsubseteq [\eta, t^\dagger]^* g$. By symmetry we obtain $t^\dagger \sqsubseteq [\eta, s^\dagger]^* h$ and therefore

$$\begin{aligned} [\eta, t^\dagger]^* g &\sqsubseteq [\eta, [\eta, s^\dagger]^* h]^* g && \text{(monotonicity)} \\ &= [\eta, s^\dagger]^* [\eta \text{ inl}, h]^* g \\ &= s^\dagger. \end{aligned}$$

We have thus shown the identity $s^\dagger = [\eta, t^\dagger]^* g$ by mutual inclusion.

- *Codiagonal.* Recall that we are claiming that

$$(T[\text{id}, \text{inr}]g)^\dagger = (g^\dagger)^\dagger$$

with $g : A \rightarrow T((B + A) + A)$. We first show that $(g^\dagger)^\dagger$ is a fixpoint of the functional defining the left-hand side as a least fixpoint, thus proving $\sqsubseteq$. That is, we have to show that

$$(g^\dagger)^\dagger = [\eta, (g^\dagger)^\dagger]^* T[\text{id}, \text{inr}]g. \quad (3.6)$$

We proceed as follows:

$$\begin{aligned}
(g^\dagger)^\dagger &= [\eta, (g^\dagger)^\dagger]^\star g^\dagger && \text{(unfolding)} \\
&= [\eta, (g^\dagger)^\dagger]^\star [\eta, g^\dagger]^\star g && \text{(unfolding)} \\
&= [[\eta, (g^\dagger)^\dagger], [\eta, (g^\dagger)^\dagger]^\star g^\dagger]^\star g \\
&= [[\eta, (g^\dagger)^\dagger], (g^\dagger)^\dagger]^\star g && \text{(unfolding)} \\
&= [[\eta, (g^\dagger)^\dagger], [\eta, (g^\dagger)^\dagger] \text{inr}]^\star g \\
&= [\eta, (g^\dagger)^\dagger]^\star T[\text{id}, \text{inr}]g.
\end{aligned}$$

For the converse inequality, we use fixpoint induction. Recall that the right hand side of the inequality at hand is a supremum of the chain $(F^i(\perp))_{i \in \mathbb{N}}$ where $F : (A \rightarrow TB) \rightarrow (A \rightarrow TB)$ is the functional defined by

$$F(f) = [\eta, f]^\star g^\dagger,$$

We show by induction on i that all members of this chain are below $(T[\text{id}, \text{inr}]g)^\dagger$, with trivial induction base. So let $f \sqsubseteq (T[\text{id}, \text{inr}]g)^\dagger$. We have to show

$$[\eta, f]^\star g^\dagger \sqsubseteq (T[\text{id}, \text{inr}]g)^\dagger.$$

We establish this by a second fixpoint induction on the definition of $g^\dagger$, again with trivial induction base. So assume that $[\eta, f]^\star h \sqsubseteq (T[\text{id}, \text{inr}]g)^\dagger$, with $h : A \rightarrow T(B + A)$; we have to show that

$$[\eta, f]^\star [\eta, h]^\star g \sqsubseteq (T[\text{id}, \text{inr}]g)^\dagger.$$

We calculate as follows:

$$\begin{aligned}
[\eta, f]^\star [\eta, h]^\star g &= [[\eta, f], [\eta, f]^\star h]^\star g \\
&\sqsubseteq [[\eta, f], (T[\text{id}, \text{inr}]g)^\dagger]^\star g && \text{(inner IH)} \\
&\sqsubseteq [[\eta, (T[\text{id}, \text{inr}]g)^\dagger], (T[\text{id}, \text{inr}]g)^\dagger]^\star g && \text{(outer IH)} \\
&= [\eta, (T[\text{id}, \text{inr}]g)^\dagger] T[\text{id}, \text{inr}]g \\
&= (T[\text{id}, \text{inr}]g)^\dagger. && \text{(unfolding)}
\end{aligned}$$

• *Uniformity.* Let $f : A \rightarrow T(X + A)$, $g : B \rightarrow T(X + B)$, $h : B \rightarrow A$ and assume that $f h = T(\text{id} + h)g$. Let us define $G(u) = [\eta, u]^\star g$, $F(u) = [\eta, u]^\star f$, and $U(u) = u h$. Then $U(\perp) = \perp$ and

$$\begin{aligned}
UF(u) &= [\eta, u]^\star f h \\
&= [\eta, u]^\star T(\text{id} + h)g \\
&= [\eta, u h]^\star g \\
&= GU(u).
\end{aligned}$$

Therefore by (3.5), $f^\dagger h = U(\mu F) = \mu G = g^\dagger$.

To prove compatibility of strength and iteration, we proceed by first showing

$$((T \text{dist})\tau(\text{id} \times f))^\dagger \sqsubseteq \tau(\text{id} \times f^\dagger).$$

First observe that, for any $g : A \rightarrow TB$,

$$\begin{array}{ccc}
 C \times (B + A) & \xrightleftharpoons[\text{dist}^{-1}]{\text{dist}} & C \times B + C \times A \\
 \text{id} \times [\eta, g] \downarrow & & \downarrow [\eta, \tau(\text{id} \times g)] \\
 C \times TB & \xrightarrow{\tau} & T(C \times B).
 \end{array} \tag{3.7}$$

This is easily checked componentwise starting from $C \times B + C \times A$ and using the fact that by definition $\text{dist}^{-1} = [\text{id} \times \text{inl}, \text{id} \times \text{inr}]$. Then we have

$$\begin{aligned}
 & \tau(\text{id} \times f^\dagger) \\
 &= \tau(\text{id} \times [\eta, f^\dagger]^* f) \\
 &= \tau(\text{id} \times [\eta, f^\dagger]^*)(\text{id} \times f) \\
 &= (\tau(\text{id} \times [\eta, f^\dagger]))^* \tau(\text{id} \times f) & (\text{STR}_4) \\
 &= ([\eta, \tau(\text{id} \times f^\dagger)] \text{dist})^* \tau(\text{id} \times f) & (3.7) \\
 &= [\eta, \tau(\text{id} \times f^\dagger)]^* (T \text{dist}) \tau(\text{id} \times f).
 \end{aligned}$$

Therefore, $\tau(\text{id} \times f^\dagger)$ is a fixed point of the functional defining $((T \text{dist}) \tau(\text{id} \times f))^\dagger$ as a least fixpoint and the inequality above holds. The converse inequality,

$$\tau(\text{id} \times f^\dagger) \sqsubseteq ((T \text{dist}) \tau(\text{id} \times f))^\dagger,$$

is shown by fixpoint induction as above for the codiagonal. The base case is trivial with

$$\tau\langle \text{fst}, \perp \text{snd} \rangle = \tau\langle \text{fst}, \perp \text{fst} \rangle = \tau\langle \text{id}, \perp \rangle \text{fst} = \perp \text{fst} = \perp.$$

Assume now that $\tau(\text{id} \times g) \sqsubseteq (T \text{dist} \tau(\text{id} \times f))^\dagger$. We can then calculate

$$\begin{aligned}
 & \tau(\text{id} \times [\eta, g]^* f) \\
 &= \tau(\text{id} \times [\eta, g]^*)(\text{id} \times f) \\
 &= (\tau(\text{id} \times [\eta, g]))^* \tau(\text{id} \times f) & (\text{STR}_4) \\
 &= ([\eta, \tau(\text{id} \times g)] \text{dist})^* \tau(\text{id} \times f) & (3.7) \\
 &\sqsubseteq ([\eta, (T \text{dist} \tau(\text{id} \times f))^\dagger] \text{dist})^* \tau(\text{id} \times f) \\
 &= [\eta, (T \text{dist} \tau(\text{id} \times f))^\dagger]^* T \text{dist} \tau(\text{id} \times f) \\
 &= (T \text{dist} \tau(\text{id} \times f))^\dagger
 \end{aligned}$$

which completes the proof. $\square$

Every complete Elgot monad $\mathbb{T}$ can express *unproductive divergence* as the generic effect

$$\perp_{X,Y} = \left(X \xrightarrow{\eta \text{ inr}} T(Y + X) \right)^\dagger : X \rightarrow TY.$$

This computation never produces any effects, i.e. behaves like a deadlock. If $\mathbb{T}$ is ω -continuous, then unproductive divergence coincides with the least element of $\text{Hom}(X, TY)$, for which reason we use the same symbol $\perp_X$, but in general, there is no ordering in which unproductive divergence could be a least element.

Lemma 3.8. *Unproductive divergence is constant, i.e. for $f : Z \rightarrow X$, we have $\perp_{X,Y} f = \perp_{Z,Y}$, and coconstant, i.e. for $h : Y \rightarrow TW$ we have $h^* \perp_{X,Y} = \perp_{X,W}$.*

Proof. Constancy: We have to show $(\eta_{Y+X} \text{inr})^\dagger f = (\eta_{Y+Z} \text{inr})^\dagger$. By uniformity, it suffices to show that $\eta_{Y+X} \text{inr} f = T(\text{id} + f) \eta_{Y+Z} \text{inr}$. We calculate the right-hand side:

$$\begin{aligned} T(\text{id} + f) \eta_{Y+Z} \text{inr} &= \eta_{Y+X} (\text{id} + f) \text{inr} && \text{(naturality of } \eta) \\ &= \eta_{Y+X} \text{inr} f. \end{aligned}$$

Coconstancy: We have

$$\begin{aligned} h^\star \perp_{X,Y} &= h^\star (\eta \text{inr}_{Y+X})^\dagger \\ &= ([T \text{inl } h, \eta_{W+X} \text{inr}]^\star \eta_{Y+X} \text{inr})^\dagger && \text{(naturality of } \dagger) \\ &= (\eta_{W+X} \text{inr})^\dagger = \perp_{X,W}. \end{aligned} \quad \square$$

The following lemma shows that there can be only one unproductive divergence:

Lemma 3.9. *Let $e : X \rightarrow T(Y + X)$ have the form $e = \eta \text{inr} u$ for $u : X \rightarrow X$. Then $e^\dagger = \perp_{X,Y}$.*

Proof. By constancy, $\perp_{X,Y} = \perp_{1,Y} !_X$, so we are to show $(\eta_{Y+1} \text{inr})^\dagger !_X = e^\dagger$. By uniformity, it suffices to show $\eta_{Y+1} \text{inr} !_X = T(\text{id} + !_X) \eta \text{inr} u$, which is immediate by naturality of η . $\square$

4. THE COINDUCTIVE GENERALIZED RESUMPTION TRANSFORMER

We proceed to recall the definition of the coinductive generalized resumption transformer [26]. We then prove our main result, stability of the class of complete Elgot monads under this construction (Theorem 5.1).

From now on we fix a strong functor $\Sigma : \mathbf{C} \rightarrow \mathbf{C}$ with strength ρ , and a strong monad $\mathbb{T}$ on $\mathbf{C}$ with strength τ . As indicated in the introduction, we think of Σ as specifying a signature of communication actions, and of T as encapsulating a notion of side-effect. We impose the following global assumption.

Assumption 4.1. *The final coalgebra of $T(X + \Sigma)$ exists for all $X \in |\mathbf{C}|$.*

We can then define a functor T_Σ whose action on objects is given by

$$T_\Sigma X = \nu \gamma. T(X + \Sigma \gamma).$$

Intuitively, $T_\Sigma X$ is a type of possibly non-terminating computation trees, in which each step triggers a computational effect specified by T , and then either terminates with a result in X or branches according to an operation from the signature represented by Σ , with arguments being again computation trees.

Remark 4.2. There are two broad classes of models satisfying Assumption 4.1:

- $\mathbf{C}$ is a locally presentable category and $\mathbb{T}$ is ranked; or
- $\mathbf{C}$ is $\omega\mathbf{Cpo}$ -enriched and has colimits of ω -chains, and T is ω -continuous (Remark 3.6).

Satisfaction of Assumption 4.1 in the first case follows from the fact that categories of coalgebras for accessible functors over locally presentable categories are again locally presentable, in particular complete [5, Exercise 2.j, Chapter 2]. This covers most of the interesting choices of base categories, such as **Set**, $\omega\mathbf{Cpo}$, various categories of predomains, and presheaf categories, as well as almost all computationally relevant monads [23, 29]. The fact that Assumption 4.1 is satisfied in the second case follows from Barr's work on algebraically

compact functors [7, Theorem 5.4], which also implies that the greatest fixed points of interest coincide with least fixed points. One example covered by the second clause but not by the first one is the continuation monad $TX = (X \rightarrow R) \rightarrow R$ on $\omega\mathbf{Cpo}$, provided that R has a least element.

Let

$$\text{out}_X : T_\Sigma X \rightarrow T(X + \Sigma T_\Sigma X)$$

be the final coalgebra structure, and let $\text{coit}(g) : Y \rightarrow T_\Sigma X$ denote the final morphism induced by a coalgebra $g : Y \rightarrow T(X + \Sigma Y)$:

$$\begin{array}{ccc} Y & \xrightarrow{\text{coit}(g)} & T_\Sigma X \\ g \downarrow & & \downarrow \text{out}_X \\ T(X + \Sigma Y) & \xrightarrow{T(X + \Sigma \text{coit}(g))} & T(X + \Sigma T_\Sigma X). \end{array}$$

Intuitively, $\text{coit}(g)$ encapsulates (in $T_\Sigma X$) a computation tree that begins by executing g , terminates in a leaf of type X if g does, and otherwise (co-)recursively continues to execute g , forming a new tree node for each recursive call. By Lambek's lemma, out_X is an isomorphism. As we see below, it is also natural in X . Thus, T maps into T_Σ via

$$\text{ext} = (T \xrightarrow{T \text{ inl}} T(\text{Id} + \Sigma T_\Sigma) \xrightarrow{\text{out}^{-1}} T_\Sigma). \quad (4.1)$$

We record explicitly that $\mathbb{T}_\Sigma$ is a strong monad:

Theorem 4.3. *Given a monad $\mathbb{T}$, T_Σ is the functorial part of a monad $\mathbb{T}_\Sigma$, with the strong monad structure denoted τ^ν , η^ν , and $(-)^*$ (for Kleisli star) and characterized by the following properties.*

- (1) *The unit $\eta^\nu : X \rightarrow T_\Sigma X$ is defined by $\text{out } \eta^\nu = \eta \text{ inl}$ (i.e. $\eta^\nu = \text{out}^{-1} \eta \text{ inl}$).*
- (2) *Given $f : X \rightarrow T_\Sigma Y$, the Kleisli lifting $f^* : T_\Sigma X \rightarrow T_\Sigma Y$ is the unique solution of the equation*

$$\text{out } f^* = [\text{out } f, \eta \text{ inr } \Sigma f^*]^* \text{out}. \quad (4.2)$$

- (3) *Given $f : X \rightarrow T_\Sigma Y$, let $g = [f, \eta^\nu] : X + Y \rightarrow T_\Sigma Y$; then g^* is a final morphism of coalgebras, namely*

$$g^* = \text{coit}([T(\text{id} + \Sigma T_\Sigma \text{ inr}) \text{out } g, \eta \text{ inr}]^* \text{out}).$$

- (4) *The strength $\tau^\nu : X \times T_\Sigma Y \rightarrow T_\Sigma(X \times Y)$ is the unique solution of*

$$\text{out } \tau^\nu = T(\text{id} + \Sigma \tau^\nu)(T\delta) \tau(\text{id} \times \text{out}) \quad (4.3)$$

with $\delta : X \times (Y + \Sigma Z) \rightarrow (X \times Y) + \Sigma(X \times Z)$ being the transformation $\delta = (\text{id} + \rho) \text{dist}$ where $\rho_{X,Y} : X \times \Sigma Y \rightarrow \Sigma(X \times Y)$ is the strength of Σ .

This justifies calling $\mathbb{T}_\Sigma$ the *coinductive generalized resumption monad* (over $\mathbb{T}$). The proof of Theorem 4.3 is facilitated by the fact that $T(X + \Sigma)$ can be shown to be a *parametrized monad*, which implies that $\mathbb{T}_\Sigma$ is a monad [34, Theorems 3.7 and 3.9]. Alternatively, the fact that $\mathbb{T}_\Sigma$ is a monad can be read off directly from the results of [26]. What is new here is that we show that $\mathbb{T}_\Sigma$ is, in fact, strong, and hence supports an interpretation of Moggi's computational metalanguage [23]. This amounts to showing that the strength defined in the last item satisfies the requisite laws in p. 3. One preliminary fact of potentially independent interest used in the proof of these laws is

Lemma 4.4. *If $T_\Sigma X$ exists for each X , then T_Σ is a functor and $\text{out} : T_\Sigma \rightarrow T(\text{Id} + \Sigma T_\Sigma)$ is a natural transformation. For any functor $G : \mathbf{B} \rightarrow \mathbf{C}$, $\text{out}_G : T_\Sigma G \rightarrow T(G + \Sigma T_\Sigma G)$ is a final $T(G + \Sigma(-))$ -coalgebra in $[\mathbf{B}, \mathbf{C}]$.*

Proof. Functoriality follows from T_Σ carrying a monad structure, as shown latter in the proof of Theorem 4.3 independently of this lemma. Now $T_\Sigma f = (\eta^\nu f)^*$, so by the description of $*$ we have

$$\begin{aligned} \text{out } T_\Sigma f &= [\text{out } \eta^\nu f, \eta \text{ inr } \Sigma T_\Sigma f]^* \text{out} \\ &= [\eta \text{ inl } f, \eta \text{ inr } \Sigma T_\Sigma f]^* \text{out} \\ &= T[\text{inl } f, \text{inr } \Sigma T_\Sigma f] \text{out} \\ &= T(f + \Sigma T_\Sigma f) \text{out}, \end{aligned}$$

i.e. out is natural.

To show finality, let $\beta : F \rightarrow T(G + \Sigma F)$ be a natural transformation. We define the universal arrow $f : F \rightarrow T_\Sigma G$ componentwise by the equation

$$\text{out } f_X = T(\text{id} + \Sigma f_X) \beta_X$$

using finality of the components $\text{out}_{GX} : T_\Sigma GX \rightarrow T(GX + \Sigma T_\Sigma GX)$. We have to show that f is natural (uniqueness is clear). So let $g : X \rightarrow Y$; we have to show $f_Y Fg = (T_\Sigma Gg) f_X$. Note that we have a $T(GY + F)$ -coalgebra

$$FX \xrightarrow{\beta_X} T(GX + \Sigma FX) \xrightarrow{T(Gg + \text{id})} T(GY + \Sigma FX);$$

we show that both $f_Y Fg$ and $(T_\Sigma Gg) f_X$ are coalgebra morphisms into $T_\Sigma GY$ for $T(Gg + \text{id}) \beta_X$. On the one hand, we have

$$\begin{aligned} \text{out } f_Y Fg &= T(\text{id} + \Sigma f_Y) \beta_Y Fg && \text{(definition of } f_Y) \\ &= T(\text{id} + \Sigma f_Y) T(Gg + \Sigma Fg) \beta_X && \text{(naturality of } \beta) \\ &= T(Gg + \Sigma(f_Y Fg)) \beta_X. \end{aligned}$$

On the other hand,

$$\begin{aligned} \text{out}(T_\Sigma Gg) f_X &= T(Gg + \Sigma T_\Sigma Gg) \text{out } f_X && \text{(naturality of out)} \\ &= T(Gg + \Sigma T_\Sigma Gg) T(\text{id} + \Sigma f_X) \beta_X && \text{(definition of } f_X) \\ &= T(Gg + \Sigma((T_\Sigma Gg) f_X)) \beta_X. \end{aligned}$$

Using the fact that there is unique morphism from a given coalgebra to the final one, we conclude that indeed $f_Y Fg = (T_\Sigma Gg) f_X$. $\square$

Proof of Theorem 4.3. Since $T(X + \Sigma)$ extends to a *parametrized monad*, T_Σ is a monad whose Kleisli lifting uniquely characterized by the corresponding equations [34, Theorems 3.7 and 3.9]. What is missing is to show that T_Σ is a strong monad, as we need here.

Let us first show the identity

$$g^* = \text{coit}([T(\text{id} + \Sigma T_\Sigma \text{ inr}) \text{out } g, \eta \text{ inr}]^* \text{out}). \quad (4.4)$$

By definition, g^* is the unique morphism making the following diagram commute:

$$\begin{array}{ccc} T_{\Sigma}(X + Y) & \xrightarrow{g^*} & T_{\Sigma}Y \\ \text{out} \downarrow & & \downarrow \text{out} \\ T(X + Y + \Sigma T_{\Sigma}(X + Y)) & \xrightarrow{[\text{out } g, \eta \text{ inr } \Sigma g^*]^*} & T(Y + \Sigma T_{\Sigma}Y) \end{array}$$

We then have on the one hand,

$$\begin{aligned} \text{out } g^* &= [\text{out}[f, \eta^\nu], \eta \text{ inr } \Sigma g^*]^* \text{ out} && (\text{definition of } _*) \\ &= [[\text{out } f, \text{out } \eta^\nu], \eta \text{ inr } \Sigma g^*]^* \text{ out} \\ &= [[\text{out } f, \eta \text{ inl}], \eta \text{ inr } \Sigma g^*]^* \text{ out} && (\text{definition of } \eta^\nu) \end{aligned}$$

and also on the other hand,

$$\begin{aligned} &T(\text{id} + \Sigma g^*) [T(\text{id} + \Sigma T_{\Sigma} \text{ inr}) \text{ out } g, \eta \text{ inr}]^* \text{ out} \\ &= [T(\text{id} + \Sigma(g^* T_{\Sigma} \text{ inr}))[\text{out } f, \text{out } \eta^\nu], \eta \text{ inr } \Sigma g^*]^* \text{ out} \\ &= [[\text{out } f, \eta \text{ inl}], \eta \text{ inr } \Sigma g^*]^* \text{ out}, \end{aligned}$$

i.e. indeed g^* satisfies the characteristic property of the final morphism (4.4).

We proceed to prove that T_{Σ} is strong. We define the strength τ^ν as the unique final coalgebra morphism shown in the following diagram:

$$\begin{array}{ccc} X \times T_{\Sigma}Y & \xrightarrow{(T\delta)\tau(\text{id} \times \text{out})} & T(X \times Y + \Sigma(X \times T_{\Sigma}Y)) \\ \tau^\nu \downarrow & & \downarrow T(\text{id} + \Sigma\tau^\nu) \\ T_{\Sigma}(X \times Y) & \xrightarrow{\text{out}} & T(X \times Y + \Sigma T_{\Sigma}(X \times Y)) \end{array}$$

That is, τ^ν is the unique solution of equation $\text{out } \tau^\nu = T(\text{id} + \Sigma\tau^\nu)(T\delta)\tau(\text{id} \times \text{out})$. By Lemma 4.4, τ^ν is a natural transformation as a composition of natural transformations. Let us check the axioms of strength from p. 3.

• (STR₁) The identity $\text{snd} = (T_{\Sigma} \text{snd})\tau^\nu$ follows from $T_{\Sigma}\langle !, \text{id} \rangle \text{snd} = \tau^\nu$ where $!$ is a suitable terminal morphism $X \rightarrow 1$, since obviously $\text{snd} = (T_{\Sigma} \text{snd})T_{\Sigma}\langle !, \text{id} \rangle \text{snd}$. Since τ^ν is uniquely defined by the corresponding characteristic identity (4.3), it suffices to show that $T_{\Sigma}\langle !, \text{id} \rangle \text{snd}$ satisfies the same identity. Indeed,

$$\begin{aligned} &T(\text{id} + \Sigma(T_{\Sigma}\langle !, \text{id} \rangle \text{snd}))(T\delta)\tau(\text{id} \times \text{out}) \\ &= T(\langle !, \text{id} \rangle \text{snd} + \Sigma(T_{\Sigma}\langle !, \text{id} \rangle \text{snd}))(T\delta)\tau(\text{id} \times \text{out}) \\ &= T(\langle !, \text{id} \rangle + \Sigma T_{\Sigma}\langle !, \text{id} \rangle)T(\text{snd} + \Sigma \text{snd})(T\delta)\tau(\text{id} \times \text{out}) \\ &= T(\langle !, \text{id} \rangle + \Sigma T_{\Sigma}\langle !, \text{id} \rangle)T(\text{snd} + \text{snd})(T \text{dist})\tau(\text{id} \times \text{out}) && (\text{STR}_1 \text{ for } \rho) \\ &= T(\langle !, \text{id} \rangle + \Sigma T_{\Sigma}\langle !, \text{id} \rangle)(T \text{snd})\tau(\text{id} \times \text{out}) && (\text{definition of dist}) \\ &= T(\langle !, \text{id} \rangle + \Sigma T_{\Sigma}\langle !, \text{id} \rangle) \text{out } \text{snd} && (\text{STR}_1 \text{ for } \tau) \\ &= \text{out}(T_{\Sigma}\langle !, \text{id} \rangle) \text{snd}. && (\text{naturality of } \tau) \end{aligned}$$

- (STR₂) In order to prove that $(T_\Sigma \text{assoc})\tau^\nu = \tau^\nu(\text{id} \times \tau^\nu) \text{assoc} : (X \times Y) \times T_\Sigma Z \rightarrow T_\Sigma((X \times Y) \times Z)$, it suffices to show that $(T_\Sigma \text{assoc}^{-1})\tau^\nu(\text{id} \times \tau^\nu) \text{assoc}$ satisfies the characteristic identity (4.3) for τ^ν , i.e.

$$\begin{aligned} & \text{out}(T_\Sigma \text{assoc}^{-1})\tau^\nu(\text{id} \times \tau^\nu) \text{assoc} \\ &= T(\text{id} + \Sigma((T_\Sigma \text{assoc}^{-1})\tau^\nu(\text{id} \times \tau^\nu) \text{assoc}))(T\delta)\tau(\text{id} \times \text{out}). \end{aligned}$$

We calculate, transforming the left hand side,

$$\begin{aligned} & \text{out}(T_\Sigma \text{assoc}^{-1})\tau^\nu(\text{id} \times \tau^\nu) \text{assoc} \\ &= T(\text{assoc}^{-1} + \Sigma T_\Sigma \text{assoc}^{-1}) \text{out} \tau^\nu(\text{id} \times \tau^\nu) \text{assoc} && \text{(naturality of out)} \\ &= T(\text{assoc}^{-1} + \Sigma T_\Sigma \text{assoc}^{-1}) \\ &\quad T(\text{id} + \Sigma \tau^\nu)(T\delta)\tau(\text{id} \times \text{out})(\text{id} \times \tau^\nu) \text{assoc} && \text{(definition of } \tau^\nu) \\ &= T(\text{assoc}^{-1} + \Sigma((T_\Sigma \text{assoc}^{-1})\tau^\nu)) \\ &\quad (T\delta)\tau(\text{id} \times T(\text{id} + \Sigma \tau^\nu)(T\delta)\tau(\text{id} \times \text{out})) \text{assoc} && \text{(definition of } \tau^\nu) \\ &= T(\text{assoc}^{-1} + \Sigma((T_\Sigma \text{assoc}^{-1})\tau^\nu)) \\ &\quad (T\delta)T(\text{id} \times (\text{id} + \Sigma \tau^\nu)\delta)\tau(\text{id} \times \tau(\text{id} \times \text{out})) \text{assoc} && \text{(naturality of } \tau). \end{aligned}$$

and then continue to transform the last part of the term:

$$\begin{aligned} & \tau(\text{id} \times \tau(\text{id} \times \text{out})) \text{assoc} \\ &= \tau(\text{id} \times \tau)(\text{id} \times (\text{id} \times \text{out})) \text{assoc} \\ &= \tau(\text{id} \times \tau) \text{assoc}((\text{id} \times \text{id}) \times \text{out}) && \text{(naturality of assoc)} \\ &= (T \text{assoc})\tau(\text{id} \times \text{out}) && \text{(STR}_2 \text{ for } \tau) \end{aligned}$$

(contracting a product of identities into an identity in the last step). Summing up, it remains to show that

$$\begin{aligned} & T(\text{assoc}^{-1} + \Sigma((T_\Sigma \text{assoc}^{-1})\tau^\nu))(T\delta)T(\text{id} \times (\text{id} + \Sigma \tau^\nu)\delta)(T \text{assoc})\tau(\text{id} \times \text{out}) \\ &= T(\text{id} + \Sigma((T_\Sigma \text{assoc}^{-1})\tau^\nu(\text{id} \times \tau^\nu) \text{assoc}))(T\delta)\tau(\text{id} \times \text{out}), \end{aligned}$$

which we reduce, removing $\tau(\text{id} \times \text{out})$ and T , multiplying from the left with $\text{assoc} + \Sigma T_\Sigma \text{assoc}$, and removing τ^ν on the left, to

$$\delta(\text{id} \times (\text{id} + \Sigma \tau^\nu)\delta) \text{assoc} = (\text{assoc} + (\Sigma(\text{id} \times \tau^\nu) \text{assoc}))\delta.$$

For the latter we calculate

$$\begin{aligned} & \delta(\text{id} \times (\text{id} + \Sigma \tau^\nu)\delta) \text{assoc} \\ &= \delta(\text{id} \times (\text{id} + \Sigma \tau^\nu))(\text{id} \times \delta) \text{assoc} \\ &= (\text{id} \times \text{id} + \Sigma(\text{id} \times \tau^\nu))\delta(\text{id} \times \delta) \text{assoc} && \text{(naturality of } \delta) \\ &= (\text{id} + \Sigma(\text{id} \times \tau^\nu))(\text{id} + \rho) \text{dist}(\text{id} \times (\text{id} + \rho) \text{dist}) \text{assoc} && \text{(definition of } \delta) \\ &= (\text{id} + \Sigma(\text{id} \times \tau^\nu))(\text{id} + \rho(\text{id} \times \rho)) \text{dist}(\text{id} \times \text{dist}) \text{assoc} && \text{(naturality of dist)} \\ &= (\text{id} + \Sigma(\text{id} \times \tau^\nu))(\text{id} + \rho(\text{id} \times \rho))(\text{assoc} + \text{assoc}) \text{dist} && (4.5) \\ &= (\text{id} + \Sigma(\text{id} \times \tau^\nu))(\text{assoc} + \Sigma \text{assoc } \rho) \text{dist} && \text{(STR}_2 \text{ for } \rho) \\ &= (\text{id} + \Sigma(\text{id} \times \tau^\nu))(\text{assoc} + \Sigma \text{assoc})\delta && \text{(definition of } \delta) \\ &= (\text{assoc} + \Sigma((\text{id} \times \tau^\nu) \text{assoc}))\delta. \end{aligned}$$

Here, we use the obvious coherence property

$$\text{dist}(\text{id} \times \text{dist}) \text{ assoc} = (\text{assoc} + \text{assoc}) \text{ dist} . \quad (4.5)$$

- (STR₃) In order to obtain the identity $\tau^\nu(\text{id} \times \eta^\nu) = \eta^\nu$, we show that the left hand side satisfies the characteristic equation for η^ν , i.e. $\text{out } \tau^\nu(\text{id} \times \eta^\nu) = \eta \text{ inl}$. Indeed,

$$\begin{aligned} \text{out } \tau^\nu(\text{id} \times \eta^\nu) &= T(\text{id} + \Sigma \tau^\nu)(T\delta)\tau(\text{id} \times \text{out})(\text{id} \times \eta^\nu) && \text{(definition of } \tau^\nu) \\ &= T(\text{id} + \Sigma \tau^\nu)(T\delta)\tau(\text{id} \times \eta)(\text{id} \times \text{inl}) && \text{(definition of } \eta^\nu) \\ &= T(\text{id} + \Sigma \tau^\nu)(T\delta)\eta(\text{id} \times \text{inl}) && \text{(STR}_3 \text{ for } \tau) \\ &= T(\text{id} + \Sigma \tau^\nu)(T\delta)T(\text{id} \times \text{inl})\eta && \text{(naturality of } \eta) \\ &= T(\text{id} + \Sigma \tau^\nu)(T \text{ inl})\eta \\ &= \eta \text{ inl} . \end{aligned}$$

- (STR₄) Given $f : X \rightarrow T_\Sigma Z$, we show that $(\tau^\nu(\text{id} \times f))^* \tau^\nu = \tau^\nu(\text{id} \times f^*)$. Let $g = [f, \eta^\nu]$ and let us show first that $(\tau^\nu(\text{id} \times g))^* \tau^\nu = \tau^\nu(\text{id} \times g^*)$. This implies the identity for f as follows:

$$\begin{aligned} (\tau^\nu(\text{id} \times f))^* \tau^\nu &= (\tau^\nu(\text{id} \times g)(\text{id} \times \text{inl}))^* \tau^\nu \\ &= (\tau^\nu(\text{id} \times g))^* T_\Sigma(\text{id} \times \text{inl}) \tau^\nu \\ &= (\tau^\nu(\text{id} \times g))^* \tau^\nu(\text{id} \times T_\Sigma \text{ inl}) && \text{(naturality of } \tau^\nu) \\ &= \tau^\nu(\text{id} \times g^*)(\text{id} \times T_\Sigma \text{ inl}) && \text{(STR}_4 \text{ for } g \text{ and } \tau^\nu) \\ &= \tau^\nu(\text{id} \times g^* T_\Sigma \text{ inl}) \\ &= \tau^\nu(\text{id} \times f^*) . \end{aligned}$$

As we have shown above both g^* and τ^ν are final morphisms from suitable coalgebras. By composing the corresponding commutative squares we obtain the following diagram:

$$\begin{array}{ccccc} X \times T_\Sigma Y & \xrightarrow{\text{id} \times [T(\text{id} + \Sigma T_\Sigma \text{ inr}) \text{ out } g, \eta \text{ inr}]^* \text{ out}} & X \times T(Z + \Sigma T_\Sigma Y) & \xrightarrow{(T\delta)\tau} & T(X \times Z + \Sigma(X \times T_\Sigma Y)) \\ \downarrow \text{id} \times g^* & & \downarrow \text{id} \times T(\text{id} + \Sigma g^*) & & \downarrow T(\text{id} + \Sigma(\tau^\nu(\text{id} \times g^*))) \\ X \times T_\Sigma Z & \xrightarrow{\text{id} \times \text{out}} & X \times T(Z + T_\Sigma Z) & \xrightarrow{(T\delta)\tau} & T(X \times Z + \Sigma(X \times T_\Sigma Z)) \\ & \searrow (T\delta)\tau(\text{id} \times \text{out}) & \downarrow T(\text{id} + \Sigma \tau^\nu) & & \\ T_\Sigma(X \times Z) & \xrightarrow{\text{out}} & T_\Sigma(X \times Z + \Sigma T_\Sigma(X \times Z)) & & \end{array}$$

from which we conclude that

$$\tau^\nu(\text{id} \times g^*) = \text{coit}((T\delta)\tau(\text{id} \times [h, \eta \text{ inr}]^* \text{ out})) .$$

where h denotes $T(\text{id} + \Sigma T_\Sigma \text{ inr}) \text{ out } g$.

We will be done once we show that also $(\tau^\nu(\text{id} \times g))^* \tau^\nu$ is a morphism from the same coalgebra to the final one, i.e. the identity

$$T(\text{id} + \Sigma((\tau^\nu(\text{id} \times g))^* \tau^\nu))(T\delta)\tau(\text{id} \times [h, \eta \text{ inr}]^* \text{out}) = \text{out}(\tau^\nu(\text{id} \times g))^* \tau^\nu. \quad (4.6)$$

Let us show that the following diagram commutes:

$$\begin{array}{ccc} T(X \times (Y + \Sigma(T_\Sigma Y))) & \xrightarrow{T\delta} & T(X \times Y + \Sigma(X \times T_\Sigma Y)) \\ (\tau(\text{id} \times [h, \eta \text{ inr}]))^* \downarrow & & \downarrow [(T\delta)\tau(\text{id} \times h), \eta \text{ inr}]^* \\ T(X \times (Z + \Sigma(T_\Sigma Y))) & \xrightarrow{T\delta} & T(X \times Z + \Sigma(X \times T_\Sigma Y)) \end{array} \quad (4.7)$$

Indeed,

$$\begin{aligned} & [(T\delta)\tau(\text{id} \times h), \eta \text{ inr}]^*(T\delta) \\ &= ([(T\delta)\tau(\text{id} \times h), \eta \text{ inr}] \delta)^* \\ &= ([(T\delta)\tau(\text{id} \times h), \eta \text{ inr}] (\text{id} + \rho) \text{dist})^* && \text{(definition of } \delta) \\ &= ([(T\delta)\tau(\text{id} \times h), \eta \text{ inr } \rho] \text{dist})^* \\ &= ([(T\delta)\tau(\text{id} \times h), \eta \delta(\text{id} \times \text{inr})] \text{dist})^* && \text{(definition of } \delta) \\ &= ([(T\delta)\tau(\text{id} \times h), (T\delta)\tau(\text{id} \times \eta)(\text{id} \times \text{inr})] \text{dist})^* && \text{(STR}_3 \text{ for } \tau) \\ &= ((T\delta)\tau[\text{id} \times h, \text{id} \times \eta \text{ inr}] \text{dist})^* \\ &= (T\delta)(\tau(\text{id} \times [h, \eta \text{ inr}]))^* \end{aligned}$$

where the last step is due to the obvious identity $[u \times v, u \times w] = (u \times [v, w]) \text{dist}^{-1}$. Finally, we obtain (4.6) as follows:

$$\begin{aligned} & T(\text{id} + \Sigma((\tau^\nu(\text{id} \times g))^* \tau^\nu)) \\ & \quad (T\delta)\tau(\text{id} \times [h, \eta \text{ inr}]^* \text{out}) \\ &= T(\text{id} + \Sigma((\tau^\nu(\text{id} \times g))^* \tau^\nu)) \\ & \quad (T\delta)(\tau(\text{id} \times [h, \eta \text{ inr}]))^* \tau(\text{id} \times \text{out}) && \text{(STR}_4 \text{ for } \tau) \\ &= T(\text{id} + \Sigma((\tau^\nu(\text{id} \times g))^* \tau^\nu)) \\ & \quad [(T\delta)\tau(\text{id} \times h), \eta \text{ inr}]^* (T\delta)\tau(\text{id} \times \text{out}) && (4.7) \\ &= T(\text{id} + \Sigma((\tau^\nu(\text{id} \times g))^* \tau^\nu)) \\ & \quad [(T\delta)\tau(\text{id} \times T(\text{id} + \Sigma T_\Sigma \text{ inr}))(\text{id} \times \text{out } g), \eta \text{ inr}]^* \\ & \quad (T\delta)\tau(\text{id} \times \text{out}) && \text{(definition of } h) \\ &= T(\text{id} + \Sigma((\tau^\nu(\text{id} \times g))^* \tau^\nu)) \\ & \quad [(T\delta)T(\text{id} \times (\text{id} + \Sigma T_\Sigma \text{ inr}))\tau(\text{id} \times \text{out } g), \eta \text{ inr}]^* \\ & \quad (T\delta)\tau(\text{id} \times \text{out}) && \text{(naturality of } \tau) \\ &= T(\text{id} + \Sigma((\tau^\nu(\text{id} \times g))^* \tau^\nu)) \\ & \quad [T(\text{id} + \Sigma(\text{id} \times T_\Sigma \text{ inr}))(T\delta)\tau(\text{id} \times \text{out } g), \eta \text{ inr}]^* \\ & \quad (T\delta)\tau(\text{id} \times \text{out}) && \text{(naturality of } \delta) \\ &= [T(\text{id} + \Sigma((\tau^\nu(\text{id} \times g))^* \tau^\nu(\text{id} \times T_\Sigma \text{ inr}))(T\delta)\tau(\text{id} \times \text{out } g), \\ & \quad \eta \text{ inr } \Sigma((\tau^\nu(\text{id} \times g))^* \tau^\nu)]^* (T\delta)\tau(\text{id} \times \text{out}) && \text{(naturality of } \eta) \end{aligned}$$

$$\begin{aligned}
 &= [T(\text{id} + \Sigma((\tau^\nu(\text{id} \times g \text{ inr}))^* \tau^\nu))(T\delta)\tau(\text{id} \times \text{out } g), \\
 &\quad \eta \text{ inr } \Sigma((\tau^\nu(\text{id} \times g))^* \tau^\nu)]^* (T\delta)\tau(\text{id} \times \text{out}) \quad (\text{naturality of } \tau^\nu) \\
 &= [T(\text{id} + \Sigma((\tau^\nu(\text{id} \times \eta^\nu))^* \tau^\nu))(T\delta)\tau(\text{id} \times \text{out } g), \\
 &\quad \eta \text{ inr } \Sigma((\tau^\nu(\text{id} \times g))^* \tau^\nu)]^* (T\delta)\tau(\text{id} \times \text{out}) \quad (\text{since } g = [f, \eta^\nu]) \\
 &= [T(\text{id} + \Sigma\tau^\nu)(T\delta)\tau(\text{id} \times \text{out } g), \\
 &\quad \eta \text{ inr } \Sigma((\tau^\nu(\text{id} \times g))^* \tau^\nu)]^* (T\delta)\tau(\text{id} \times \text{out}) \quad (\text{STR}_3 \text{ for } \tau^\nu) \\
 &= [T(\text{id} + \Sigma\tau^\nu)(T\delta)\tau(\text{id} \times \text{out})(\text{id} \times g), \\
 &\quad \eta \text{ inr } \Sigma((\tau^\nu(\text{id} \times g))^* \tau^\nu)]^* (T\delta)\tau(\text{id} \times \text{out}) \\
 &= [\text{out } \tau^\nu(\text{id} \times g), \eta \text{ inr } \Sigma((\tau^\nu(\text{id} \times g))^* \tau^\nu)]^* (T\delta)\tau(\text{id} \times \text{out}) \quad (\text{definition of } \tau^\nu) \\
 &= [\text{out } \tau^\nu(\text{id} \times g), \eta \text{ inr } \Sigma(\tau^\nu(\text{id} \times g))^*]^* \\
 &\quad T(\text{id} + \Sigma\tau^\nu)(T\delta)\tau(\text{id} \times \text{out}) \\
 &= [\text{out } \tau^\nu(\text{id} \times g), \eta \text{ inr } \Sigma(\tau^\nu(\text{id} \times g))^*]^* \text{out } \tau^\nu \quad (\text{definition of } \tau^\nu) \\
 &= \text{out}(\tau^\nu(\text{id} \times g))^* \tau^\nu. \quad (\text{definition of } _*)
 \end{aligned}$$

We have thus shown all properties (STR₁)–(STR₄) and the proof is completed. $\square$

Following Uustalu [34] (and other work [26, 1]), we next introduce a notion of guardedness.

Definition 4.5. (Guardedness) A morphism $f : X \rightarrow T_\Sigma(Y + Z)$ is *guarded* if there is $u : X \rightarrow T(Y + \Sigma T_\Sigma(Y + Z))$ such that $\text{out } f = T(\text{inl} + \text{id}) u$:

$$\begin{array}{ccc}
 X & \xrightarrow{f} & T_\Sigma(Y + Z) \\
 u \downarrow & & \downarrow \text{out} \\
 T(Y + \Sigma T_\Sigma(Y + Z)) & \xrightarrow{T(\text{inl} + \text{id})} & T((Y + Z) + \Sigma T_\Sigma(Y + Z)).
 \end{array}$$

Guardedness of $f : X \rightarrow T_\Sigma(Y + Z)$ intuitively means that any call to a computation of type Z in f occurs only under a free operation, i.e. via the right hand summand in $T((Y + Z) + \Sigma T_\Sigma(Y + Z))$. A familiar instance of this notion occurs in process algebra [9], illustrated in simplified form as follows.

Example 4.6. Let $\mathbb{T}$ be the countable powerset monad over a suitable category, i.e. $\mathbb{T}X = \mathcal{P}_{\omega_1} X = \{Y \subseteq X \mid |Y| \leq \omega\}$. Take $\Sigma = A \times (-)$; then the object $T_\Sigma X = \nu\gamma. \mathcal{P}_{\omega_1}(X + A \times \gamma)$ can be considered as the domain of possibly infinite countably nondeterministic processes over actions from A with final results in X . A morphism $n \rightarrow T_\Sigma(X + n)$ can be seen as a system of n mutually recursive process definitions; the latter is guarded in the sense of Definition 4.5 iff every recursive call of a process is preceded by an action, which coincides with the standard notion of guardedness from process algebra. We recall an example of an *unguarded* definition in this setting in Section 8.

5. ITERATION ON COINDUCTIVE RESUMPTIONS

We next establish one of the main technical contributions of the paper by proving that iteration operators, i.e. Elgot monad structures, propagate uniquely along extensions $\mathbb{T} \rightarrow \mathbb{T}_\Sigma$.

Theorem 5.1. *Let $\mathbb{T}$ be a complete Elgot monad and let $\mathbb{T}_\Sigma$ be the monad identified in Theorem 4.3, i.e. the coinductive generalized resumption monad over $\mathbb{T}$.*

- (1) *There is a unique iteration operator making $\mathbb{T}_\Sigma$ a complete Elgot monad that extends iteration in $\mathbb{T}$ in the sense that for $f : X \rightarrow T_\Sigma(Y + X)$ and $g : X \rightarrow T(Y + X)$, if*

$$\text{out } f = (T \text{ inl}) g \quad (5.1)$$

(i.e. $f = \text{out}^{-1}(T \text{ inl}) g$) then

$$\text{out } f^\dagger = (T \text{ inl}) g^\dagger. \quad (5.2)$$

- (2) *For any guarded morphism $f : X \rightarrow T_\Sigma(Y + X)$, $f^\dagger$ is the unique morphism satisfying the unfolding property $[\eta^\nu, f^\dagger]^* f = f^\dagger$.*

The proof of Theorem 5.1 relies on a fairly complicated chain of calculations and will, to aid readability, be partitioned into separate lemmas. Before we dive into these details, let us outline the general idea.

Uustalu already proves that guarded morphisms f have unique iterates $f^\dagger$ [34, Theorem 3.11], which readily implies the second clause. In showing the first clause of Theorem 5.1, the key step is then to define $f^\dagger$ for unrestricted f in a consistent manner. For $f : X \rightarrow T_\Sigma(Y + X)$, let $\star f : X \rightarrow T_\Sigma(Y + X)$ be the composite

$$\begin{aligned} X &\xrightarrow{w^\dagger} T(Y + \Sigma T_\Sigma(Y + X)) \\ &\xrightarrow{T(\text{inl} + \text{id})} T((Y + X) + \Sigma T_\Sigma(Y + X)) \\ &\xrightarrow{\text{out}^{-1}} T_\Sigma(Y + X) \end{aligned}$$

(guarded by definition), where w is the composite

$$\begin{aligned} X &\xrightarrow{f} T_\Sigma(Y + X) \\ &\xrightarrow{\text{out}} T((Y + X) + \Sigma T_\Sigma(Y + X)) \\ &\xrightarrow{T\pi} T((Y + \Sigma T_\Sigma(Y + X)) + X) \end{aligned}$$

with $\pi = [\text{inl} + \text{id}, \text{inl inr}]$. That is, $\star f$ makes f guarded by iterating

$$\text{out } f : X \rightarrow T((Y + X) + \Sigma T_\Sigma(Y + X))$$

(in the complete Elgot monad $\mathbb{T}$) over the middle summand X . It is easy to check that $\star f = f$ when f is guarded. We hence can consistently define

$$f^\dagger = (\star f)^\dagger \quad (5.3)$$

(in $\mathbb{T}_\Sigma$). The remaining technical challenge is now to prove that this definition indeed satisfies the axioms of complete Elgot monads and that it is the unique such iteration operator on $\mathbb{T}_\Sigma$ extending the given iteration operator on $\mathbb{T}$.

Lemma 5.2. *Given $f : X \rightarrow T_\Sigma(Y + X)$, $f^\dagger : f : X \rightarrow T_\Sigma Y$ defined by (5.3) satisfies unfolding, naturality, dinaturality and uniformity.*

Proof. To make sure that definition (5.3) introduces the iteration consistently with the iteration for guarded morphisms we check that $\star f = f$ whenever f is guarded. Suppose that $\text{out } f = T(\text{inl} + \text{id})u$. Then $f = \text{out}^{-1} T(\text{inl} + \text{id})u$ and therefore

$$\begin{aligned} \star f &= \text{out}^{-1} T(\text{inl} + \text{id})((T\pi) \text{out } f)^\dagger \\ &= \text{out}^{-1} T(\text{inl} + \text{id})((T\pi) \text{out } \text{out}^{-1} T(\text{inl} + \text{id})u)^\dagger \\ &= \text{out}^{-1} T(\text{inl} + \text{id})((T\pi)T(\text{inl} + \text{id})u)^\dagger \\ &= \text{out}^{-1} T(\text{inl} + \text{id})(T \text{inl } u)^\dagger \\ &= \text{out}^{-1} T(\text{inl} + \text{id})u \\ &= f. \end{aligned}$$

Let us check *unfolding*, *naturality*, *dinaturality* and *uniformity* (Definition 3.1) in order.

- *Unfolding.* For any $f : X \rightarrow T_\Sigma(Y + X)$ we have

$$\begin{aligned} f^\dagger &= [\eta^\nu, f^\dagger]^\star \star f && \text{(definition of } _^\dagger) \\ &= [\eta^\nu, f^\dagger]^\star \text{out}^{-1} T(\text{inl} + \text{id})((T\pi) \text{out } f)^\dagger && \text{(definition of } \star) \\ &= \text{out}^{-1} [[\eta \text{inl}, \text{out } f^\dagger], \eta \text{inr } \Sigma[\eta^\nu, f^\dagger]^\star]^\star \\ &\quad T(\text{inl} + \text{id})((T\pi) \text{out } f)^\dagger && \text{(definition of } _^\star) \\ &= \text{out}^{-1} [\eta \text{inl}, \eta \text{inr } \Sigma[\eta^\nu, f^\dagger]^\star]^\star ((T\pi) \text{out } f)^\dagger \\ &= \text{out}^{-1} T(\text{id} + \Sigma[\eta^\nu, f^\dagger]^\star)((T\pi) \text{out } f)^\dagger \end{aligned}$$

and thus we obtain the following intermediate equation:

$$\text{out } f^\dagger = T(\text{id} + \Sigma[\eta^\nu, f^\dagger]^\star)((T\pi) \text{out } f)^\dagger. \quad (5.4)$$

Now, continuing the above calculation we obtain

$$\begin{aligned} f^\dagger &= \text{out}^{-1} T(\text{id} + \Sigma[\eta^\nu, f^\dagger]^\star)((T\pi) \text{out } f)^\dagger \\ &= \text{out}^{-1} T(\text{id} + \Sigma[\eta^\nu, f^\dagger]^\star)[\eta, ((T\pi) \text{out } f)^\dagger]^\star (T\pi) \text{out } f && \text{(unfolding)} \\ &= \text{out}^{-1} [T(\text{id} + \Sigma[\eta^\nu, f^\dagger]^\star)\eta, \text{out } f^\dagger]^\star (T\pi) \text{out } f && (5.4) \\ &= \text{out}^{-1} [\eta(\text{id} + \Sigma[\eta^\nu, f^\dagger]^\star), \text{out } f^\dagger]^\star (T\pi) \text{out } f && \text{(naturality of } \eta) \\ &= \text{out}^{-1} [[\eta \text{inl}, \text{out } f^\dagger], \eta \text{inr } \Sigma[\eta^\nu, f^\dagger]^\star]^\star \text{out } f \\ &= \text{out}^{-1} [\text{out}[\eta^\nu, f^\dagger], \eta \text{inr } \Sigma[\eta^\nu, f^\dagger]^\star]^\star \text{out } f && \text{(definition of } \eta^\nu) \\ &= [\eta^\nu, f^\dagger]^\star \text{out}^{-1} \text{out } f && \text{(naturality of } _^\star) \\ &= [\eta^\nu, f^\dagger]^\star f. \end{aligned}$$

- *Naturality.* Assume that $h : X \rightarrow T_\Sigma(Y + X)$ is guarded and show that so is $[(T_\Sigma \text{inl})g, \eta^\nu \text{inr}]^\star h$ for any $g : Y \rightarrow Z$. Let u be such that $\text{out } h = T(\text{inl} + \text{id})u$ and let

$w = [(T_\Sigma \text{ inl})g, \eta^\nu \text{ inr}]$. Then

$$\begin{aligned}
& \text{out}[(T_\Sigma \text{ inl})g, \eta^\nu \text{ inr}]^* h \\
&= [\text{out } w, \eta \text{ inr } \Sigma w^*]^* \text{out } h \\
&= [\text{out } w, \eta \text{ inr } \Sigma w^*]^* T(\text{inl} + \text{id})u \\
&= [\text{out } w \text{ inl}, \eta \text{ inr } \Sigma w^*]^* u \\
&= [\text{out}(T_\Sigma \text{ inl})g, \eta \text{ inr } \Sigma w^*]^* u \\
&= [T(\text{inl} + \Sigma T_\Sigma \text{ inl}) \text{out } g, \eta \text{ inr } \Sigma w^*]^* u \\
&= T(\text{inl} + \text{id})[T(\text{id} + \Sigma T_\Sigma \text{ inl}) \text{out } g, \eta \text{ inr } \Sigma w^*]^* u.
\end{aligned}$$

Now, since $t = [(T_\Sigma \text{ inl})g, \eta^\nu \text{ inr}]^* \star f$ is guarded, it is the unique fixpoint of the map

$$t \mapsto [\eta^\nu, t]^* [(T_\Sigma \text{ inl})g, \eta^\nu \text{ inr}]^* \star f.$$

However, on the other hand,

$$\begin{aligned}
& [\eta^\nu, g^* f^\dagger]^* [(T_\Sigma \text{ inl})g, \eta^\nu \text{ inr}]^* \star f \\
&= [g, g^* f^\dagger]^* \star f \\
&= [g, g^* (\star f)^\dagger]^* \star f \\
&= g^* [\eta^\nu, (\star f)^\dagger]^* \star f \\
&= g^* f^\dagger
\end{aligned}$$

and therefore $t^\dagger = g^* f^\dagger$. It remains to show that

$$[(T_\Sigma \text{ inl})g, \eta^\nu \text{ inr}]^* \star f = \star [(T_\Sigma \text{ inl})g, \eta^\nu \text{ inr}]^* f.$$

Since $\star f$ is guarded by definition, we know by the calculation above that $[(T_\Sigma \text{ inl})g, \eta^\nu \text{ inr}]^* \star f$ is guarded and therefore

$$[(T_\Sigma \text{ inl})g, \eta \text{ inr}]^* \star f = \star [(T_\Sigma \text{ inl})g, \eta \text{ inr}]^* f.$$

To finish the proof, we calculate

$$\star [(T_\Sigma \text{ inl})g, \eta^\nu \text{ inr}]^* \star f = \text{out}^{-1} T(\text{inl} + \text{id})((T\pi) \text{out}[(T_\Sigma \text{ inl})g, \eta^\nu \text{ inr}]^* \star f)^\dagger.$$

Further transforming the dagger expression in the previous term yields

$$\begin{aligned}
& ((T\pi) \text{out}[(T_\Sigma \text{ inl})g, \eta \text{ inr}]^* \star f)^\dagger \\
&= ((T\pi)[\text{out } w, \eta \text{ inr } \Sigma w^*]^* T(\text{inl} + \text{id})((T\pi) \text{out } f)^\dagger)^\dagger \\
&= ((T\pi)[\text{out}(T_\Sigma \text{ inl})g, \eta \text{ inr } \Sigma w^*]^* ((T\pi) \text{out } f)^\dagger)^\dagger \\
&= (((T \text{ inl})(T\pi)[\text{out}(T_\Sigma \text{ inl})g, \eta \text{ inr } \Sigma w^*], \eta \text{ inr})^* (T\pi) \text{out } f)^\dagger)^\dagger \quad (\text{nat. for } \mathbb{T}) \\
&= (T[\text{id}, \text{inr}]((T \text{ inl})(T\pi)[\text{out}(T_\Sigma \text{ inl})g, \eta \text{ inr } \Sigma w^*], \eta \text{ inr})^* \\
&\quad (T\pi) \text{out } f)^\dagger \quad (\text{codiag. for } \mathbb{T}) \\
&= (((T\pi) \text{out}(T_\Sigma \text{ inl})g, \eta \text{ inr}], (T\pi) \eta \text{ inr } \Sigma w^*)^* \text{out } f)^\dagger \\
&= ((T\pi)[[\text{out}(T_\Sigma \text{ inl})g, \eta \text{ inl inr}], \eta \text{ inr } \Sigma w^*]^* \text{out } f)^\dagger \\
&= ((T\pi)[\text{out}[(T_\Sigma \text{ inl})g, \eta^\nu \text{ inr}], \eta \text{ inr } \Sigma w^*]^* \text{out } f)^\dagger \\
&= ((T\pi) \text{out}[(T_\Sigma \text{ inl})g, \eta^\nu \text{ inr}]^* f)^\dagger
\end{aligned}$$

and therefore

$$\star[(T_\Sigma \text{ inl})g, \eta \text{ inr}]^* \star f = \star[(T_\Sigma \text{ inl})g, \eta \text{ inr}]^* f.$$

• *Dinaturality.* Given $g : X \rightarrow T_\Sigma(Y + Z)$ and $h : Z \rightarrow T_\Sigma(Y + X)$, let $s = [\eta^\nu \text{ inl}, h]^* g : X \rightarrow T_\Sigma(Y + X)$, $t = [\eta^\nu \text{ inl}, g]^* h : Z \rightarrow T_\Sigma(Y + Z)$, $w = [\eta^\nu, t^\dagger]^* g : X \rightarrow T_\Sigma Y$. The idea is to show the identity

$$[\eta^\nu, w]^* \star s = [\eta^\nu, t^\dagger]^* [\eta^\nu \text{ inl}, \star t]^* g \quad (5.5)$$

from which we will be able to obtain that

$$\begin{aligned} w &= [\eta^\nu, t^\dagger]^* g \\ &= [\eta^\nu, [\eta^\nu, t^\dagger]^* \star t]^* g && \text{(unfolding)} \\ &= [\eta^\nu, t^\dagger]^* [\eta^\nu \text{ inl}, \star t]^* g \\ &= [\eta^\nu, w]^* \star s, \end{aligned} \quad (5.5)$$

i.e. that w satisfies the recursive equation uniquely identifying $s^\dagger$ and therefore we obtain the desired equation $w = s^\dagger$. Let us introduce the following morphisms:

$$\begin{aligned} \mathbf{l} &= T((\text{id} + \Sigma[\eta^\nu \text{ inl}, h]^*) + \text{id}) : T(Y + T_\Sigma(Y + Z) + Z) \rightarrow T(Y + T_\Sigma(Y + X) + Z), \\ \mathbf{r} &= T((\text{id} + \Sigma[\eta^\nu \text{ inl}, g]^*) + \text{id}) : T(Y + T_\Sigma(Y + X) + X) \rightarrow T(Y + T_\Sigma(Y + Z) + X) \end{aligned}$$

and observe that

$$\begin{aligned} ((T\pi) \text{ out } s)^\dagger &= ((T\pi) [\text{out}[\eta^\nu \text{ inl}, h], \eta \text{ inr } \Sigma[\eta^\nu \text{ inl}, h]^*]^\star \text{ out } g)^\dagger \\ &= ([\eta \text{ inl}(\text{id} + \Sigma[\eta^\nu \text{ inl}, h]^*), (T\pi) \text{ out } h]^\star (T\pi) \text{ out } g)^\dagger \\ &= ([\eta \text{ inl}, (T\pi) \text{ out } h]^\star \mathbf{l}(T\pi) \text{ out } g)^\dagger. \end{aligned}$$

An analogous calculation applies to $((T\pi) \text{ out } t)^\dagger$ and hence we obtain

$$((T\pi) \text{ out } s)^\dagger = ([\eta \text{ inl}, (T\pi) \text{ out } h]^\star \mathbf{l}(T\pi) \text{ out } g)^\dagger, \quad (5.6)$$

$$((T\pi) \text{ out } t)^\dagger = ([\eta \text{ inl}, (T\pi) \text{ out } g]^\star \mathbf{r}(T\pi) \text{ out } h)^\dagger. \quad (5.7)$$

Let us calculate the left-hand side of (5.5):

$$\begin{aligned} &\text{out}[\eta^\nu, w]^* \star s \\ &= [\text{out}[\eta^\nu, w], \eta \text{ inr } \Sigma[\eta^\nu, w]^*]^\star \text{out } \star s \\ &= [\text{out}[\eta^\nu, w], \eta \text{ inr } \Sigma[\eta^\nu, w]^*]^\star T(\text{inl} + \text{id})((T\pi) \text{ out } s)^\dagger \\ &= T(\text{id} + \Sigma[\eta^\nu, w]^*)((T\pi) \text{ out } s)^\dagger \\ &= T(\text{id} + \Sigma[\eta^\nu, w]^*)([\eta \text{ inl}, (T\pi) \text{ out } h]^\star \mathbf{l}(T\pi) \text{ out } g)^\dagger && (5.6) \\ &= T(\text{id} + \Sigma[\eta^\nu, w]^*) \\ &\quad [\eta, ([\eta \text{ inl}, \mathbf{l}(T\pi) \text{ out } g]^\star (T\pi) \text{ out } h)^\dagger]^\star \mathbf{l}(T\pi) \text{ out } g \\ &= T(\text{id} + \Sigma[\eta^\nu, w]^*)[\eta(\text{id} + \Sigma[\eta^\nu \text{ inl}, h]^*), \\ &\quad ([\eta \text{ inl}, \mathbf{l}(T\pi) \text{ out } g]^\star (T\pi) \text{ out } h)^\dagger]^\star (T\pi) \text{ out } g \\ &= [\eta(\text{id} + \Sigma[\eta^\nu, t^\dagger]^*), \end{aligned}$$

$$T(\text{id} + \Sigma[\eta^\nu, w]^*)([\eta \text{ inl}, \mathbf{l}(T\pi) \text{ out } g]^*(T\pi) \text{ out } h)^\dagger]^*(T\pi) \text{ out } g,$$

where for the last step, we called on the following calculation:

$$\begin{aligned} & [\eta^\nu, w]^*[\eta^\nu \text{ inl}, h]^* \\ &= [\eta^\nu, [\eta^\nu, w]^*h]^* \\ &= [\eta^\nu, [\eta^\nu, [\eta^\nu, t^\dagger]^*g]^*h]^* \\ &= [\eta^\nu, [\eta^\nu, t^\dagger]^*[\eta^\nu \text{ inl}, g]^*h]^* \\ &= [\eta^\nu, t^\dagger]^*. \end{aligned}$$

Now, let us calculate the right-hand side of (5.5):

$$\begin{aligned} & \text{out}[\eta^\nu, t^\dagger]^*[\eta^\nu \text{ inl}, \star t]^*g \\ &= [\text{out}[\eta^\nu, t^\dagger], \eta \text{ inr } \Sigma[\eta^\nu, t^\dagger]^*]^* \text{out}[\eta^\nu \text{ inl}, \star t]^*g \\ &= [\text{out}[\eta^\nu, t^\dagger], \eta \text{ inr } \Sigma[\eta^\nu, t^\dagger]^*]^* [\text{out}[\eta^\nu \text{ inl}, \star t], \eta \text{ inr } \Sigma[\eta^\nu \text{ inl}, \star t]^*]^* \text{out } g \\ &= [\text{out}[\eta^\nu, t^\dagger], \eta \text{ inr } \Sigma[\eta^\nu, t^\dagger]^*]^* [\eta(\text{id} + \Sigma[\eta^\nu \text{ inl}, \star t]^*), \text{out } \star t]^*(T\pi) \text{ out } g \\ &= [\eta(\text{id} + \Sigma[\eta^\nu, t^\dagger]^*), [\text{out}[\eta^\nu, t^\dagger], \eta \text{ inr } \Sigma[\eta^\nu, t^\dagger]^*]^*]^* \text{out } \star t]^*(T\pi) \text{ out } g. \end{aligned}$$

We have thus reduced (5.5) to

$$\begin{aligned} & T(\text{id} + \Sigma[\eta^\nu, w]^*)([\eta \text{ inl}, \mathbf{l}((T\pi) \text{ out } g)]^*(T\pi) \text{ out } h)^\dagger \\ &= [\text{out}[\eta^\nu, t^\dagger], \eta \text{ inr } \Sigma[\eta^\nu, t^\dagger]^*]^* \text{out } \star t. \end{aligned}$$

Then, on the one hand

$$\begin{aligned} & T(\text{id} + \Sigma[\eta^\nu, w]^*)([\eta \text{ inl}, \mathbf{l}(T\pi) \text{ out } g]^*((T\pi) \text{ out } h))^\dagger \\ &= (T((\text{id} + \Sigma[\eta^\nu, w]^*) + \text{id})[\eta \text{ inl}, \mathbf{l}(T\pi) \text{ out } g]^*((T\pi) \text{ out } h))^\dagger \\ &= ([\eta \text{ inl}(\text{id} + \Sigma[\eta^\nu, w]^*), T((\text{inl} + \Sigma[\eta^\nu, w]^*) + \text{id})\mathbf{l}(T\pi) \text{ out } g]^*(T\pi) \text{ out } h)^\dagger \\ &= ([\eta \text{ inl}(\text{id} + \Sigma[\eta^\nu, w]^*), T((\text{inl} + \Sigma[\eta^\nu, t^\dagger]^*) + \text{id})(T\pi) \text{ out } g]^*(T\pi) \text{ out } h)^\dagger \\ &= (T((\text{id} + \Sigma[\eta^\nu, t^\dagger]^*) + \text{id})[\eta \text{ inl}(\text{id} + \Sigma[\eta^\nu \text{ inl}, g]^*), (T\pi) \text{ out } g]^*(T\pi) \text{ out } h)^\dagger \\ &= T(\text{id} + \Sigma[\eta^\nu, t^\dagger]^*)([\eta \text{ inl}(\text{id} + \Sigma[\eta^\nu \text{ inl}, g]^*), (T\pi) \text{ out } g]^*(T\pi) \text{ out } h)^\dagger \\ &= T(\text{id} + \Sigma[\eta^\nu, t^\dagger]^*)([\eta \text{ inl}, (T\pi) \text{ out } g]^*\mathbf{r}(T\pi) \text{ out } h)^\dagger \end{aligned}$$

and also on the other hand

$$\begin{aligned} & [\text{out}[\eta^\nu, t^\dagger], \eta \text{ inr } \Sigma[\eta^\nu, t^\dagger]^*]^* \text{out } \star t \\ &= [\text{out}[\eta^\nu, t^\dagger], \eta \text{ inr } \Sigma[\eta^\nu, t^\dagger]^*]^* T(\text{inl} + \text{id})((T\pi) \text{ out } t)^\dagger \\ &= T(\text{id} + \Sigma[\eta^\nu, t^\dagger]^*)((T\pi) \text{ out } t)^\dagger \\ &= T(\text{id} + \Sigma[\eta^\nu, t^\dagger]^*)([\eta \text{ inl}, (T\pi) \text{ out } g]^*\mathbf{r}(T\pi) \text{ out } h)^\dagger. \end{aligned} \tag{5.7}$$

We have thus reduced both sides of (5.5) to identical expression.

- *Uniformity.* First, show uniformity under the assumption that g is guarded. Suppose that $fh = T_\Sigma(\text{id} + h)g$. It is then sufficient to verify that $f^\dagger h$ satisfies the unfolding identity for g . Indeed,

$$\begin{aligned} f^\dagger h &= [\eta^\nu, f^\dagger]^* fh \\ &= [\eta^\nu, f^\dagger]^* T_\Sigma(\text{id} + h)g \\ &= [\eta^\nu, f^\dagger h]^* g. \end{aligned}$$

Now consider the general case. Suppose that again we have $fh = T_\Sigma(\text{id} + h)g$. We prove the following auxiliary identity:

$$((T\pi) \text{ out } f)^\dagger h = T(\text{id} + \Sigma T_\Sigma(\text{id} + h))((T\pi) \text{ out } g)^\dagger. \quad (5.8)$$

Observe that

$$\begin{aligned} ((T\pi) \text{ out } f)h &= (T\pi) \text{ out } T_\Sigma(\text{id} + h)g \\ &= (T\pi)T(\text{id} + h + \Sigma T_\Sigma(\text{id} + h)) \text{ out } g \\ &= T(\text{id} + h)T((\text{id} + \Sigma T_\Sigma(\text{id} + h)) + \text{id})(T\pi) \text{ out } g, \end{aligned}$$

from which by uniformity of the iteration operator of $\mathbb{T}$, we obtain

$$((T\pi) \text{ out } f)^\dagger h = (T((\text{id} + \Sigma T_\Sigma(\text{id} + h)) + \text{id})(T\pi) \text{ out } g)^\dagger.$$

After transforming the right hand side by naturality of the iteration operator of $\mathbb{T}$ we arrive at (5.8).

Next we prove that $(\star f)h = T_\Sigma(\text{id} + h)\star g$:

$$\begin{aligned} (\star f)h &= \text{out}^{-1} T(\text{inl} + \text{id})((T\pi) \text{ out } f)^\dagger h && \text{(definition of } \star) \\ &= \text{out}^{-1} T(\text{inl} + \text{id})T(\text{id} + \Sigma T_\Sigma(\text{id} + h))((T\pi) \text{ out } g)^\dagger && (5.8) \\ &= \text{out}^{-1} T((\text{id} + h) + \Sigma T_\Sigma(\text{id} + h))T(\text{inl} + \text{id})((T\pi) \text{ out } g)^\dagger \\ &= T_\Sigma(\text{id} + h) \text{out}^{-1} T(\text{inl} + \text{id})((T\pi) \text{ out } g)^\dagger && \text{(Lemma 4.4)} \\ &= T_\Sigma(\text{id} + h)\star g. && \text{(definition of } \star) \end{aligned}$$

We have shown before that for guarded g uniformity holds, and therefore $f^\dagger h = (\star f)^\dagger h = (\star g)^\dagger = g^\dagger$. $\square$

We now deal with the last axiom, *codiagonal*, whose proof is more involved than that of the other properties and therefore handled in a separate lemma:

Lemma 5.3. *The assignment of $f^\dagger : X \rightarrow T_\Sigma Y$ to $f : X \rightarrow T_\Sigma(Y + X)$ defined by (5.3) satisfies the codiagonal axiom.*

Proof. Let $g : X \rightarrow T_\Sigma((Y + X) + X)$. We shall show below that

$$\star(T_\Sigma[\text{id}, \text{inr}]\star g) = \star(T_\Sigma[\text{id}, \text{inr}]g). \quad (5.9)$$

Since $T_\Sigma[\text{id}, \text{inr}]^\dagger g$ is the unique fixpoint of the map

$$\gamma \mapsto [\eta^\nu, \gamma]^* \star(T_\Sigma[\text{id}, \text{inr}]g)$$

we will be done once we show that $(g^\dagger)^\dagger$ is also a fixpoint of the same map, i.e.

$$(g^\dagger)^\dagger = [\eta^\nu, (g^\dagger)^\dagger]^* \star(T_\Sigma[\text{id}, \text{inr}]g). \quad (5.10)$$

Let us again denote by $\pi : (Y + X) + X \rightarrow (Y + X) + X$ the morphism swapping the last two components of the coproduct. We consider the following three cases.

(1) $T_\Sigma[\text{id}, \text{inr}]g$ is guarded. Then we obtain (5.10) directly as follows

$$\begin{aligned}
(g^\dagger)^\dagger &= [\eta^\nu, (g^\dagger)^\dagger]^\ast g^\dagger && \text{(unfolding)} \\
&= [\eta^\nu, (g^\dagger)^\dagger]^\ast [\eta^\nu, g^\dagger]^\ast g && \text{(unfolding)} \\
&= [[\eta^\nu, (g^\dagger)^\dagger], [\eta^\nu, (g^\dagger)^\dagger]^\ast g^\dagger]^\ast g \\
&= [[\eta^\nu, (g^\dagger)^\dagger], (g^\dagger)^\dagger]^\ast g && \text{(unfolding)} \\
&= [\eta^\nu, (g^\dagger)^\dagger]^\ast T_\Sigma[\text{id}, \text{inr}]g \\
&= [\eta^\nu, (g^\dagger)^\dagger]^\ast \star(T_\Sigma[\text{id}, \text{inr}]g).
\end{aligned}$$

(2) $(T_\Sigma\pi)g$ is guarded. E.g. let $(T_\Sigma\pi)g = \text{out}^{-1}T(\text{inl} + \text{id})u$. Then $T_\Sigma[\text{id}, \text{inr}] \star g$ is also guarded, which is certified by the following calculation, involving the definitions of g , $\star$ and the naturality axiom for $_^\dagger$:

$$\begin{aligned}
&T_\Sigma[\text{id}, \text{inr}] \star g \\
&= T_\Sigma[\text{id}, \text{inr}] \star ((T_\Sigma\pi) \text{out}^{-1}T(\text{inl} + \text{id})u) \\
&= T_\Sigma[\text{id}, \text{inr}] \text{out}^{-1}T(\text{inl} + \text{id}) \left((T\pi) \text{out}(T_\Sigma\pi) \text{out}^{-1}T(\text{inl} + \text{id})u \right)^\dagger \\
&= T_\Sigma[\text{id}, \text{inr}] \text{out}^{-1}T(\text{inl} + \text{id}) \left((T\pi)T(\pi + \Sigma T_\Sigma\pi)T(\text{inl} + \text{id})u \right)^\dagger \\
&= T_\Sigma[\text{id}, \text{inr}] \text{out}^{-1}T(\text{inl} + \text{id}) \left(T((\text{inl} + \text{id}) + \text{id})(T\pi)T(\text{id} + \Sigma T_\Sigma\pi)u \right)^\dagger \\
&= T_\Sigma[\text{id}, \text{inr}] \text{out}^{-1}T(\text{inl} + \text{id})T(\text{inl} + \text{id}) \left((T\pi)T(\text{id} + \Sigma T_\Sigma\pi)u \right)^\dagger \\
&= T_\Sigma[\text{id}, \text{inr}] \text{out}^{-1}T(\text{inl} \text{inl} + \text{id}) \left((T\pi)T(\text{id} + \Sigma T_\Sigma\pi)u \right)^\dagger \\
&= \text{out}^{-1}T([\text{id}, \text{inr}] + \Sigma T_\Sigma[\text{id}, \text{inr}])T(\text{inl} \text{inl} + \text{id}) \left((T\pi)T(\text{id} + \Sigma T_\Sigma\pi)u \right)^\dagger \\
&= \text{out}^{-1}T(\text{inl} + \Sigma T_\Sigma[\text{id}, \text{inr}]) \left((T\pi)T(\text{id} + \Sigma T_\Sigma\pi)u \right)^\dagger \\
&= \text{out}^{-1}T(\text{inl} + \text{id})T(\text{id} + \Sigma T_\Sigma[\text{id}, \text{inr}]) \left((T\pi)T(\text{id} + \Sigma T_\Sigma\pi)u \right)^\dagger.
\end{aligned}$$

The proof of (5.10) now can be completed as follows:

$$\begin{aligned}
(g^\dagger)^\dagger &= ((\star g)^\dagger)^\dagger && \text{(definition of } _^\dagger) \\
&= [\eta^\nu, ((\star g)^\dagger)^\dagger]^\ast \star(T_\Sigma[\text{id}, \text{inr}] \star g) && \text{(Clause (1))} \\
&= [\eta^\nu, (g^\dagger)^\dagger]^\ast \star(T_\Sigma[\text{id}, \text{inr}]g). && (5.9)
\end{aligned}$$

(3) g is guarded. Let $h = (T_\Sigma\pi) \star (T_\Sigma\pi)g$. It is easy to see that h is guarded. We use the following identity

$$\star g^\dagger = [\eta^\nu, g^\dagger]^\ast h \quad (5.11)$$

whose proof runs as follows. Let $g = \text{out}^{-1}T(\text{inl} + \text{id})u$ for some u and observe that $\pi \text{inl} = (\text{inl} + \text{id})$. We apply out to the right-hand side of the equation,

$$\begin{aligned}
 & \text{out}[\eta^\nu, g^\dagger]^*(T_\Sigma\pi) \star (T_\Sigma\pi)g \\
 &= [\text{out}[\eta^\nu, g^\dagger], \eta \text{inr} \Sigma[\eta^\nu, g^\dagger]^*]^\star \\
 & \quad \text{out}(T_\Sigma\pi) \text{out}^{-1}T(\text{inl} + \text{id})((T\pi) \text{out}(T_\Sigma\pi)g)^\dagger \quad (\text{defn. of } _^\dagger, \star) \\
 &= [\text{out}[\eta^\nu, g^\dagger]\pi \text{inl}, \eta \text{inr} \Sigma([\eta^\nu, g^\dagger]^*T_\Sigma\pi)]^\star ((T\pi) \text{out}(T_\Sigma\pi)g)^\dagger \\
 &= [\text{out}[\eta^\nu \text{inl}, g^\dagger], \eta \text{inr} \Sigma([\eta^\nu, g^\dagger]^*T_\Sigma\pi)]^\star \\
 & \quad ((T\pi)T(\pi + \Sigma T_\Sigma\pi) \text{out} g)^\dagger \\
 &= ([[(T \text{inl})\text{out}[\eta^\nu \text{inl}, g^\dagger], \eta \text{inr}([\eta^\nu, g^\dagger]^*\Sigma T_\Sigma\pi)], \eta \text{inr}]^\star \\
 & \quad (T\pi)T(\pi + \Sigma T_\Sigma\pi) \text{out} g)^\dagger \quad (\text{naturality}) \\
 &= ([[(T \text{inl}) \text{out}[\eta^\nu \text{inl}, g^\dagger], \eta \text{inl} \text{inr} \Sigma([\eta^\nu, g^\dagger]^*T_\Sigma\pi)], \eta \text{inr}]^\star \\
 & \quad (T\pi)T(\pi + \Sigma T_\Sigma\pi) \text{out} g)^\dagger \\
 &= ([[(T \text{inl}) \text{out}[\eta^\nu \text{inl}, g^\dagger], \eta \text{inr}], \eta \text{inl} \text{inr} \Sigma([\eta^\nu, g^\dagger]^*T_\Sigma\pi)]^\star \\
 & \quad T(\pi + \Sigma T_\Sigma\pi) \text{out} g)^\dagger \quad (\text{defn. } T\pi) \\
 &= ([[(T \text{inl}) \text{out}[\eta^\nu \text{inl}, g^\dagger], \eta \text{inr}]\pi, \\
 & \quad \eta \text{inl} \text{inr} \Sigma([\eta^\nu, g^\dagger]^*(T_\Sigma\pi)(T_\Sigma\pi))]^\star T(\text{inl} + \text{id})u)^\dagger \quad (g \text{ guarded}) \\
 &= ([[(T \text{inl}) \text{out}[\eta^\nu \text{inl}, g^\dagger], \eta \text{inr}]\pi \text{inl}, \eta \text{inl} \text{inr} \Sigma[\eta^\nu, g^\dagger]^*]^\star u)^\dagger \\
 &= ([[\eta \text{inl} \text{inl} \text{inl}, \eta \text{inr}], \eta \text{inl} \text{inr} \Sigma[\eta^\nu, g^\dagger]^*]^\star u)^\dagger \\
 &= ([[\eta(\text{inl} \text{inl} + \text{id}), \eta \text{inl} \text{inr} \Sigma[\eta^\nu, g^\dagger]^*]^\star u)^\dagger.
 \end{aligned}$$

On the other hand, applying out to the left-hand side yields the same result:

$$\begin{aligned}
 & \text{out} \star (g^\dagger) \\
 &= T(\text{inl} + \text{id})((T\pi) \text{out}(g^\dagger))^\dagger \\
 &= ([[(T \text{inl})\eta(\text{inl} + \text{id}), \eta \text{inr}]^\star (T\pi) \text{out}(g^\dagger))^\dagger \quad (\text{naturality}) \\
 &= ([[\eta \text{inl} \text{inl} \text{inl}, \text{inl} \text{inr}], \eta \text{inr}]^\star (T\pi) \text{out}[\eta^\nu, g^\dagger]^*g)^\dagger \\
 &= ([[\eta(\text{inl} \text{inl} + \text{id}), \eta \text{inl} \text{inr}]^\star [\text{out}[\eta^\nu, g^\dagger], \eta \text{inr} \Sigma[\eta^\nu, g^\dagger]^*]^\star \text{out} g)^\dagger \quad (\text{defn. } T\pi) \\
 &= ([[\eta(\text{inl} \text{inl} + \text{id}), \eta \text{inl} \text{inr}]^\star [[\eta \text{inl}, \text{out} g^\dagger], \eta \text{inr} \Sigma[\eta^\nu, g^\dagger]^*]^\star \\
 & \quad T(\text{inl} + \text{id})u)^\dagger \quad (g \text{ guarded}) \\
 &= ([[\eta(\text{inl} \text{inl} + \text{id}), \eta \text{inl} \text{inr}]^\star [\eta \text{inl}, \eta \text{inr} \Sigma[\eta^\nu, g^\dagger]^*]^\star u)^\dagger \\
 &= ([[\eta(\text{inl} \text{inl} + \text{id}), \eta \text{inl} \text{inr} \Sigma[\eta^\nu, g^\dagger]^*]^\star u)^\dagger.
 \end{aligned}$$

Then the goal can be obtained as follows. First, observe the following:

$$\begin{aligned}
 (g^\dagger)^\dagger &= ([\eta^\nu, g^\dagger]^*h)^\dagger \quad (5.11) \\
 &= ([[\eta^\nu \text{inl}, \eta^\nu \text{inr}], g^\dagger]^*h)^\dagger \\
 &= ([[\eta^\nu \text{inl}, g^\dagger], \eta \text{inr}]^\star \star (T_\Sigma\pi) g)^\dagger \quad (\text{defn. of } \pi)
 \end{aligned}$$

$$\begin{aligned}
&= ((T_\Sigma[\text{id}, \text{inr}])([\eta^\nu \text{ inl inl}, T_\Sigma \text{ inl } g^\dagger], \eta \text{ inr})^* \star (T_\Sigma \pi) g)^\dagger \\
&= ((([\eta^\nu \text{ inl inl}, T_\Sigma \text{ inl } g^\dagger], \eta \text{ inr})^* \star (T_\Sigma \pi) g)^\dagger)^\dagger && \text{(Clause (2))} \\
&= (((T_\Sigma \text{ inl}[\eta^\nu \text{ inl}, g^\dagger], \eta \text{ inr})^* \star (T_\Sigma \pi) g)^\dagger)^\dagger \\
&= ([\eta^\nu \text{ inl}, g^\dagger]^* \star (T_\Sigma \pi) g)^\dagger^\dagger && \text{(naturality)} \\
&= ([\eta^\nu \text{ inl}, g^\dagger]^* ((T_\Sigma \pi) g)^\dagger)^\dagger && \text{(defn. of } _^\dagger) \\
&= [\eta^\nu, ([\eta^\nu \text{ inl}, g^\dagger]^* ((T_\Sigma \pi) g)^\dagger)^\dagger]^* [\eta^\nu \text{ inl}, g^\dagger]^* ((T_\Sigma \pi) g)^\dagger && \text{(unfolding)} \\
&= [\eta^\nu, (g^\dagger)^\dagger]^* [\eta^\nu \text{ inl}, g^\dagger]^* ((T_\Sigma \pi) g)^\dagger \\
&= [\eta^\nu, [\eta^\nu, (g^\dagger)^\dagger]^* g^\dagger]^* ((T_\Sigma \pi) g)^\dagger \\
&= [\eta^\nu, (g^\dagger)^\dagger]^* ((T_\Sigma \pi) g)^\dagger && \text{(unfolding)}
\end{aligned}$$

It is easy to see that $((T_\Sigma \pi) g)^\dagger$ is guarded, and hence, by the previous calculation, $(g^\dagger)^\dagger = (((T_\Sigma \pi) g)^\dagger)^\dagger$. Finally, by Clause (2), $((T_\Sigma \pi) g)^\dagger = ((T_\Sigma[\text{id}, \text{inr}])(T_\Sigma \pi) g)^\dagger = (T_\Sigma[\text{id}, \text{inr}]g)^\dagger$.
(4) g is unrestricted. Then,

$$\begin{aligned}
(g^\dagger)^\dagger &= ((\star g)^\dagger)^\dagger \\
&= ((T_\Sigma[\text{id}, \text{inr}]) \star g)^\dagger && \text{(Clause (3))} \\
&= (\star (T_\Sigma[\text{id}, \text{inr}]) \star (T_\Sigma \pi) g)^\dagger \\
&= (\star (T_\Sigma[\text{id}, \text{inr}]) (T_\Sigma \pi) g)^\dagger && (5.9) \\
&= (T_\Sigma[\text{id}, \text{inr}]g)^\dagger
\end{aligned}$$

and we are done. It remains to prove (5.9). Observe that by definition,

$$\begin{aligned}
&\star (T_\Sigma[\text{id}, \text{inr}]) \star (T_\Sigma \pi) g \\
&= \star T_\Sigma[\text{id}, \text{inr}] \text{out}^{-1} T(\text{inl} + \text{id})((T\pi) \text{out } g)^\dagger \\
&= \text{out}^{-1} T(\text{inl} + \text{id})((T\pi) \text{out } T_\Sigma[\text{id}, \text{inr}] \text{out}^{-1} T(\text{inl} + \text{id})((T\pi) \text{out } g)^\dagger)^\dagger
\end{aligned}$$

Let us further transform the expression after $\text{out}^{-1} T(\text{inl} + \text{id})$:

$$\begin{aligned}
&((T\pi) T([\text{id}, \text{inr}] + \Sigma T_\Sigma[\text{id}, \text{inr}]) T(\text{inl} + \text{id})((T\pi) \text{out } g)^\dagger)^\dagger \\
&= ((T\pi) T(\text{id} + \Sigma T_\Sigma[\text{id}, \text{inr}])(T\pi) \text{out } g)^\dagger^\dagger \\
&= (([T \text{ inl } \pi \eta(\text{id} + \Sigma T_\Sigma[\text{id}, \text{inr}]), \eta \text{ inr}]^* (T\pi) \text{out } g)^\dagger)^\dagger && \text{(naturality)} \\
&= (([T\pi] \eta(\text{id} + \Sigma T_\Sigma[\text{id}, \text{inr}]), (T\pi) \eta \text{ inl inr}]^* (T\pi) \text{out } g)^\dagger && \text{(codiagonal)} \\
&= ((T\pi)[[\eta \text{ inl}, \eta \text{ inl inr}], \eta \text{ inr } \Sigma T_\Sigma[\text{id}, \text{inr}]]^* \text{out } g)^\dagger \\
&= ((T\pi)[\text{out}(\eta^\nu[\text{id}, \text{inr}]), \eta \text{ inr } \Sigma T_\Sigma[\text{id}, \text{inr}]]^* \text{out } g)^\dagger \\
&= ((T\pi) \text{out}(T_\Sigma[\text{id}, \text{inr}]g)^\dagger)^\dagger.
\end{aligned}$$

Therefore,

$$\begin{aligned}
&\star (T_\Sigma[\text{id}, \text{inr}]) \star (T_\Sigma \pi) g \\
&= \text{out}^{-1} T(\text{inl} + \text{id})((T\pi) \text{out}(T_\Sigma[\text{id}, \text{inr}]g)^\dagger)^\dagger \\
&= \star T_\Sigma[\text{id}, \text{inr}]g
\end{aligned}$$

and we are done. $\square$

Lemma 5.4. *The assignment of $f^\dagger : X \rightarrow T_\Sigma Y$ to $f : X \rightarrow T_\Sigma(Y + X)$ defined by (5.3) is compatible with strength, i.e.*

$$\tau^\nu(\text{id} \times f^\dagger) = ((T_\Sigma \text{dist})\tau^\nu(\text{id} \times f))^\dagger.$$

Proof. Let f be guarded with $\text{out } f = T(\text{inl} + \text{id})u$. Then, $f' = (T_\Sigma \text{dist})\tau^\nu(\text{id} \times f)$ is also guarded with $\text{out } f' = T(\text{inl} + \text{id})T(\text{id} + \Sigma((T_\Sigma \text{dist})\tau^\nu))(T\delta)\tau(\text{id} \times u)$ where δ is as in Theorem 4.3 (besides guardedness of f , the proof of this equation uses naturality of out and the definitions of τ and dist). The following calculation shows that $\tau^\nu(\text{id} \times f^\dagger)$ satisfies the unfolding property for $((T_\Sigma \text{dist})\tau^\nu(\text{id} \times f))^\dagger$:

$$\begin{aligned} & \tau^\nu(\text{id} \times f^\dagger) \\ &= \tau^\nu(\text{id} \times [\eta^\nu, f^\dagger]^* f) \\ &= \tau^\nu(\text{id} \times [\eta^\nu, f^\dagger]^*)(\text{id} \times f) \\ &= (\tau^\nu(\text{id} \times [\eta^\nu, f^\dagger]))^* \tau^\nu(\text{id} \times f) & (\text{STR}_4) \\ &= ([\eta^\nu, \tau^\nu(\text{id} \times f^\dagger)] \text{dist})^* \tau^\nu(\text{id} \times f) & (3.7) \\ &= [\eta^\nu, \tau^\nu(\text{id} \times f^\dagger)]^* (T_\Sigma \text{dist})\tau^\nu(\text{id} \times f) \\ &= [\eta^\nu, \tau^\nu(\text{id} \times f^\dagger)]^* \blackstar (T_\Sigma \text{dist})\tau^\nu(\text{id} \times f), \end{aligned}$$

and hence $\tau^\nu(\text{id} \times f^\dagger)$ and $((T_\Sigma \text{dist})\tau^\nu(\text{id} \times f))^\dagger$ are equal.

The general case reduces to the guarded case by means of the equation

$$(T_\Sigma \text{dist})\tau^\nu(\text{id} \times \blackstar f) = \blackstar ((T_\Sigma \text{dist})\tau^\nu(\text{id} \times f)), \quad (5.12)$$

as follows:

$$\begin{aligned} \tau^\nu(\text{id} \times f^\dagger) &= \tau^\nu(\text{id} \times (\blackstar f)^\dagger) & (\text{definition of } _^\dagger) \\ &= ((T \text{dist})\tau^\nu(\text{id} \times \blackstar f))^\dagger \\ &= (\blackstar (T_\Sigma \text{dist} \tau^\nu(\text{id} \times f)))^\dagger & (5.12) \\ &= (T_\Sigma \text{dist} \tau^\nu(\text{id} \times f))^\dagger. & (\text{definition of } _^\dagger) \end{aligned}$$

We show (5.12) by establishing commutativity of the following diagram where $Q = C \times B + C \times A$ (the identity in question is read from the border):

$$\begin{array}{ccc}
& C \times A & \\
\text{id} \times ((T\pi) \text{out } f)^\dagger \swarrow & & \searrow ((T\pi) \text{out } (T_\Sigma \text{dist}) \tau^\nu (\text{id} \times f))^\dagger \\
C \times T(B + \Sigma T_\Sigma(B + A)) & \xrightarrow{T(\text{id} + \Sigma T_\Sigma \text{dist}) T(\text{id} + \Sigma \tau^\nu) (T\delta) \tau} & T(C \times B + \Sigma T_\Sigma Q) \\
\downarrow \text{id} \times T(\text{inl} + \text{id}) & & \downarrow T(\text{inl} + \text{id}) \\
C \times T((B + A) + \Sigma T_\Sigma(B + A)) & \xrightarrow{T(\text{dist} + \Sigma T_\Sigma \text{dist}) T(\text{id} + \Sigma \tau^\nu) (T\delta) \tau} & T(Q + \Sigma T_\Sigma Q) \\
& \searrow (T_\Sigma \text{dist}) \tau^\nu (\text{id} \times \text{out}^{-1}) & \swarrow \text{out}^{-1} \\
& T_\Sigma Q &
\end{array}$$

The bottom triangle commutes as follows:

$$\begin{aligned}
& (T_\Sigma \text{dist}) \tau^\nu (\text{id} \times \text{out}^{-1}) \\
&= (T_\Sigma \text{dist}) \text{out}^{-1} T(\text{id} + \Sigma \tau^\nu) (T\delta) \tau (\text{id} \times \text{out}) (\text{id} \times \text{out}^{-1}) \\
&= \text{out}^{-1} \text{out} (T_\Sigma \text{dist}) \text{out}^{-1} T(\text{id} + \Sigma \tau^\nu) (T\delta) \tau \\
&= \text{out}^{-1} T(\text{dist} + \Sigma T_\Sigma \text{dist}) T(\text{id} + \Sigma \tau^\nu) (T\delta) \tau.
\end{aligned}$$

The middle square commutes by properties of τ , dist and δ :

$$\begin{aligned}
& T(\text{dist} + \Sigma(T_\Sigma \text{dist } \tau^\nu))(T\delta) \tau (\text{id} \times T(\text{inl} + \text{id})) \\
&= T(\text{dist} + \Sigma(T_\Sigma \text{dist } \tau^\nu))(T\delta) T(\text{id} \times (\text{inl} + \text{id})) \tau && \text{(naturality of } \tau) \\
&= T(\text{dist} + \Sigma(T_\Sigma \text{dist } \tau^\nu)) T((\text{id} \times \text{inl}) + \text{id}) (T\delta) \tau && \text{(naturality of } \delta) \\
&= T(\text{dist}(\text{id} \times \text{inl}) + \text{id}) T(\text{id} + \Sigma(T_\Sigma \text{dist } \tau^\nu)) (T\delta) \tau \\
&= T(\text{inl} + \text{id}) T(\text{id} + \Sigma(T_\Sigma \text{dist } \tau^\nu)) (T\delta) \tau.
\end{aligned}$$

This leaves us with the top triangle. Let $\alpha = (\text{id} + \Sigma(T_\Sigma \text{dist } \tau^\nu))\delta$. We apply the assumption that τ is compatible with iteration to $(T\alpha)\tau(\text{id} \times ((T\pi) \text{out } f)^\dagger)$ and further calculate as follows:

$$\begin{aligned}
& (T\alpha)\tau(\text{id} \times ((T\pi) \text{out } f)^\dagger) \\
&= (T\alpha)(T \text{dist } \tau(\text{id} \times (T\pi) \text{out } f))^\dagger \\
&= (T(\alpha + \text{id}))(T \text{dist}) \tau(\text{id} \times (T\pi) \text{out } f))^\dagger && \text{(naturality)} \\
&= (T(\alpha + \text{id}))(T \text{dist}) T(\text{id} \times \pi) \tau(\text{id} \times \text{out})(\text{id} \times f))^\dagger && \text{(naturality of } \tau) \\
&= (T((\text{id} + \Sigma(T_\Sigma \text{dist } \tau^\nu) + \text{id})) \\
&\quad T(\delta + \text{id})(T \text{dist}) T(\text{id} \times \pi) \tau(\text{id} \times \text{out})(\text{id} \times f))^\dagger && \text{(definition of } \alpha) \\
&= (T((\text{id} + \Sigma(T_\Sigma \text{dist } \tau^\nu) + \text{id})) \\
&\quad T((\text{id} + \rho) + \text{id}) T(\text{dist} + \text{id})(T \text{dist}) T(\text{id} \times \pi) \\
&\quad \tau(\text{id} \times \text{out})(\text{id} \times f))^\dagger. && \text{(definition of } \delta)
\end{aligned}$$

At this position we apply the obvious identity

$$T(\text{dist} + \text{id})(T \text{ dist})T(\text{id} \times \pi) = (T\pi)T(\text{dist} + \text{id})(T \text{ dist})$$

and then proceed as follows:

$$\begin{aligned}
 & (T((\text{id} + \Sigma(T_\Sigma \text{ dist } \tau^\nu) + \text{id})) \\
 & \quad T((\text{id} + \rho) + \text{id})(T\pi)T(\text{dist} + \text{id})(T \text{ dist}) \\
 & \quad \tau(\text{id} \times \text{out})(\text{id} \times f))^\dagger \\
 = & (T((\text{id} + \Sigma(T_\Sigma \text{ dist } \tau^\nu) + \text{id})) \\
 & \quad (T\pi)T(\text{dist} + \rho)(T \text{ dist})\tau(\text{id} \times \text{out})(\text{id} \times f))^\dagger \\
 = & (T((\text{id} + \Sigma(T_\Sigma \text{ dist } \tau^\nu)) + \text{id})) \\
 & \quad (T\pi)T(\text{dist} + \text{id})(T\delta)\tau(\text{id} \times \text{out})(\text{id} \times f))^\dagger \quad (\text{definition of } \delta) \\
 = & ((T\pi)T(\text{dist} + \Sigma T_\Sigma \text{ dist})) \\
 & \quad T(\text{id} + \Sigma \tau^\nu)(T\delta)\tau(\text{id} \times \text{out})(\text{id} \times f))^\dagger \\
 = & ((T\pi)T(\text{dist} + \Sigma T_\Sigma \text{ dist}) \text{ out } \tau^\nu(\text{id} \times f))^\dagger \quad (\text{definition of } \tau^\nu) \\
 = & ((T\pi) \text{ out}(T_\Sigma \text{ dist}) \tau^\nu(\text{id} \times f))^\dagger.
 \end{aligned}$$

This yields the proof of the top triangle of the diagram and therefore completes the proof of the lemma. $\square$

Finally, we can return to the proof of Theorem 5.1.

Proof of Theorem 5.1. As we indicated above, the second clause is already proved by Uustalu [34]. To show the existence part of the first clause we call on the above Lemmas 5.2, 5.3 and 5.4 and additionally prove that iteration on $\mathbb{T}_\Sigma$ extends iteration on $\mathbb{T}$, i.e. that (5.1) implies (5.2). Let us call morphism f for which there is g satisfying (5.1) *completely unguarded*. Suppose that (5.1) holds. Then the proof of (5.2) runs as follows:

$$\begin{aligned}
 \text{out } f^\dagger &= (\star f)^\dagger \\
 &= \text{out}(\text{out}^{-1}T(\text{inl} + \text{id})((T\pi) \text{ out } f)^\dagger)^\dagger \\
 &= \text{out}(\text{out}^{-1}T(\text{inl} + \text{id})((T\pi)(T \text{ inl})g)^\dagger)^\dagger \quad (5.1) \\
 &= \text{out}(\text{out}^{-1}T(\text{inl} + \text{id})((T \text{ inl} + \text{id})g)^\dagger)^\dagger \\
 &= \text{out}(\text{out}^{-1}T(\text{inl} + \text{id})(T \text{ inl})g^\dagger)^\dagger \quad (\text{naturality}) \\
 &= \text{out}(\text{out}^{-1}T(\text{inl inl})g^\dagger)^\dagger \\
 &= \text{out}[\eta^\nu, f^\dagger]^\star \text{out}^{-1}T(\text{inl inl})g^\dagger \quad (\text{unfolding}) \\
 &= \text{out}[\eta^\nu, f^\dagger]^\star \text{out}^{-1}T(\text{inl inl})g^\dagger \\
 &= [\text{out}[\eta^\nu, f^\dagger], \eta \text{ inr } \Sigma[\eta^\nu, f^\dagger]^\star]^\star T(\text{inl inl})g^\dagger \quad (4.2) \\
 &= [\text{out } \eta^\nu]^\star g^\dagger \\
 &= [\eta \text{ inl}]^\star g^\dagger \\
 &= (T \text{ inl})g^\dagger.
 \end{aligned}$$

It remains to show the uniqueness part of the first clause. To that end we first show that any morphism $f : X \rightarrow T_\Sigma(Y + X)$ can be decomposed by means of morphisms $g : X \rightarrow T_\Sigma(Z + X)$ and $h : Z \rightarrow T_\Sigma(Y + X)$, where $Z = Y + \Sigma T_\Sigma(Y + X)$, as

$$f = [h, \eta^\nu \text{inr}]^* g \quad (5.13)$$

with completely unguarded g . Next we show that

$$f^\dagger = (h^* g^\dagger)^\dagger \quad (5.14)$$

and that

$$h^* g^\dagger = \clubsuit f. \quad (5.15)$$

In summary, we obtain that $f^\dagger = (h^* g^\dagger)^\dagger = (\clubsuit f)^\dagger$. The following proofs of (5.14) and (5.15) do not depend on the concrete definition of $_^\dagger$ on $\mathbb{T}_\Sigma$ but only use its abstract properties as an iteration operator of a complete Elgot monad and compatibility with the underlying iteration operator for $\mathbb{T}$. Hence, the identity $f^\dagger = (\clubsuit f)^\dagger$ would be valid for any other such operator, but since $(\clubsuit f)^\dagger$ is uniquely defined all of them must agree.

Let $g = \text{out}^{-1} T(\text{inl } \pi) \text{out } f$ (recall that $\pi = [\text{inl} + \text{id}, \text{inl inr}]$), which is, by definition, completely unguarded, and let $h = \text{out}^{-1} \eta(\text{inl} + \text{id})$.

Then the proof of (5.13) runs as follows:

$$\begin{aligned} & [h, \eta^\nu \text{inr}]^* g \\ &= [\text{out}^{-1} \eta(\text{inl} + \text{id}), \eta^\nu \text{inr}]^* g \\ &= [\text{out}^{-1} \eta(\text{inl} + \text{id}), \text{out}^{-1} \eta \text{inl inr}]^* g && \text{(Theorem 4.3)} \\ &= (\text{out}^{-1} \eta[\text{inl} + \text{id}, \text{inl inr}])^* g \\ &= (\text{out}^{-1} \eta \pi)^* g \\ &= \text{out}^{-1} [\eta \pi, \eta \text{inr } \Sigma(\text{out}^{-1} \eta \pi)^*]^\star \text{out } g && \text{(Theorem 4.3)} \\ &= \text{out}^{-1} (\eta [\pi, \text{inr } \Sigma(\text{out}^{-1} \eta \pi)^*])^\star T(\text{inl } \pi) \text{out } f \\ &= \text{out}^{-1} (\eta \pi)^\star (T \pi) \text{out } f \\ &= \text{out}^{-1} (T \pi) (T \pi) \text{out } f \\ &= f. \end{aligned}$$

Next, we show (5.14):

$$\begin{aligned} (h^* g^\dagger)^\dagger &= ((T_\Sigma \text{inl } h, \eta^\nu \text{inr}]^* g)^\dagger)^\dagger && \text{(naturality)} \\ &= (T_\Sigma [\text{id}, \text{inr}] [(T_\Sigma \text{inl } h, \eta^\nu \text{inr}]^* g)^\dagger)^\dagger && \text{(codiagonal)} \\ &= ([h, T_\Sigma [\text{id}, \text{inr}] \eta^\nu \text{inr}]^* g)^\dagger \\ &= ([h, \eta^\nu \text{inr}]^* g)^\dagger \\ &= f^\dagger. \end{aligned}$$

Finally, we prove (5.15):

$$\begin{aligned}
 h^*g^\dagger &= (\text{out}^{-1}\eta(\text{inl} + \text{id}))^*g^\dagger && (\text{definition of } _*) \\
 &= \text{out}^{-1}[\eta(\text{inl} + \text{id}), \eta \text{inr } \Sigma h^*]^* \text{out } g^\dagger \\
 &= \text{out}^{-1}[\eta(\text{inl} + \text{id}), \eta \text{inr } \Sigma h^*]^*(T\text{inl})((T\pi) \text{out } f)^\dagger && (g \text{ compl. ung.}) \\
 &= \text{out}^{-1}T(\text{inl} + \text{id})((T\pi) \text{out } f)^\dagger \\
 &= \star f.
 \end{aligned}$$

This finishes the proof. $\square$

6. A COPRODUCT CHARACTERIZATION OF COINDUCTIVE RESUMPTIONS

Our second main result is a universal characterization of the coinductive resumption monad transformer; essentially, we show that T_Σ arises as the coproduct of $\mathbb{T}$ with the free complete Elgot monad over Σ (see Section 9 for discussion of a similar result on *completely iterative* monads). This result relies on a natural notion of morphism between complete Elgot monads:

Definition 6.1. A *complete Elgot monad morphism* $\xi : \mathbb{R} \rightarrow \mathbb{S}$ between complete Elgot monads $\mathbb{R}, \mathbb{S}$ is a morphism ξ between the underlying strong monads (i.e. $\xi\eta = \eta$, $\xi f^* = (\xi f)^*\xi$ for $f : X \rightarrow RY$, and $\xi\tau = \tau(\text{id} \times \xi)$, see [22]) additionally satisfying

$$(\xi g)^\dagger = \xi g^\dagger$$

for $g : X \rightarrow R(Y + X)$. Complete Elgot monads over $\mathbf{C}$ and their morphisms form an (overlarge) category $\mathbf{CElg}(\mathbf{C})$. We have a forgetful functor from $\mathbf{CElg}(\mathbf{C})$ to the category of strong functors and strong natural transformations; mention of *free* complete Elgot monads refers to this forgetful functor.

Note next that the coinductive resumption monad T_Σ implements, by construction, all the operations of Σ , that is, we have a canonical strong natural transformation $\iota^\mathbb{T} : \Sigma \rightarrow T_\Sigma$, given by

$$\iota_X^\mathbb{T} = \text{out}^{-1}\eta \text{inr } \Sigma\eta^\nu$$

where the typing of the composite is shown in

$$\Sigma X \xrightarrow{\Sigma\eta^\nu} ST_\Sigma X \xrightarrow{\text{inr}} X + ST_\Sigma X \xrightarrow{\eta} T(X + ST_\Sigma X) \xrightarrow{\text{out}^{-1}} T_\Sigma X.$$

Moreover, recall that T maps into T_Σ via a natural transformation $\text{ext} : T \rightarrow T_\Sigma$ defined in Equation (4.1). We have

Lemma 6.2. *The natural transformation $\text{ext} : \mathbb{T} \rightarrow \mathbb{T}_\Sigma$ is a complete Elgot monad morphism.*

Proof. Let us verify the identities

$$\xi\eta = \eta \quad \xi f^* = (\xi f)^*\xi \quad \xi\tau = \tau(\text{id} \times \xi) \quad (\xi g)^\dagger = \xi g^\dagger \quad (6.1)$$

with $f : X \rightarrow TY$ and $g : X \rightarrow T(Y + X)$ from left to right.

- Compatibility of ext with unit is a straightforward consequence of Theorem 4.3: $\text{ext } \eta = \text{out}^{-1}(T\text{inl})\eta = \text{out}^{-1}\eta \text{inl} = \eta^\nu$.

- In order to show compatibility of ext with Kleisli star we call the definition of the latter from Theorem 4.3:

$$\begin{aligned}
(\text{ext } g)^* \text{ext} &= (\text{out}^{-1}(T\text{inl})g)^* \text{out}^{-1}(T\text{inl}) \\
&= \text{out}^{-1}[\text{out } \text{out}^{-1}(T\text{inl})g, \eta \text{inr } \Sigma(\text{ext } g)^*]^*(T\text{inl}) \\
&= \text{out}^{-1}((T\text{inl})g)^* \\
&= \text{out}^{-1}(T\text{inl})g^* \\
&= \text{ext } g^*.
\end{aligned}$$

- Recall the distributivity transformation $\delta : A \times (B + \Sigma C) \rightarrow A \times B + \Sigma(A \times C)$ from Theorem 4.3. Then by the corresponding definition of τ^ν ,

$$\begin{aligned}
\tau^\nu(\text{id} \times \text{ext}) &= \text{out}^{-1}T(\text{id} + \Sigma\tau^\nu)(T\delta)\tau(\text{id} \times \text{out ext}) \\
&= \text{out}^{-1}T(\text{id} + \Sigma\tau^\nu)(T\delta)\tau(\text{id} \times T\text{inl}) \\
&= \text{out}^{-1}T(\text{id} + \Sigma\tau^\nu)T(\text{id} + \rho)(T\text{dist})T(\text{id} + \text{inl})\tau \\
&= \text{out}^{-1}T(\text{id} + \Sigma\tau^\nu)T(\text{id} + \rho)(T\text{inl})\tau \\
&= \text{out}^{-1}T(\text{id} + \Sigma\tau^\nu)(T\text{inl})\tau \\
&= \text{out}^{-1}(T\text{inl})\tau \\
&= \text{ext } \tau.
\end{aligned}$$

- Since $\text{out ext } g = (T_\Sigma \text{inl})g$, then by Theorem 5.1, $\text{out}(\text{ext } g)^\dagger = (T\text{inl})g^\dagger$, from which the last identity in (6.1) follows by composition with out^{-1} on the left. $\square$

Summing up, we have, slightly abusing notation, a cospan of strong natural transformations

$$\mathbb{T} \xrightarrow{\text{ext}} \mathbb{T}_\Sigma \xleftarrow{\iota^\mathbb{T}} \Sigma$$

with the left arrow being a complete Elgot monad morphism. It turns out that this gives a universal characterization of $\mathbb{T}_\Sigma$:

Theorem 6.3. *The cospan $\mathbb{T} \xrightarrow{\text{ext}} \mathbb{T}_\Sigma \xleftarrow{\iota^\mathbb{T}} \Sigma$ is universal. Explicitly: Given a complete Elgot monad $\mathbb{S}$, a strong natural transformation $v : \Sigma \rightarrow \mathbb{S}$, and a complete Elgot monad morphism $\sigma : \mathbb{T} \rightarrow \mathbb{S}$, there exists a unique complete Elgot monad morphism $\xi : \mathbb{T}_\Sigma \rightarrow \mathbb{S}$ such that $\xi \text{ext} = \sigma$ and $\xi \iota^\mathbb{T} = v$:*

$$\begin{array}{ccc}
\mathbb{T} & \xrightarrow{\text{ext}} & \mathbb{T}_\Sigma \xleftarrow{\iota^\mathbb{T}} \Sigma \\
& \searrow \sigma & \downarrow \xi \quad \swarrow v \\
& & \mathbb{S}.
\end{array} \tag{6.2}$$

Specifically, ξ is given as $\xi = \zeta^\dagger$ with ζ defined componentwise by

$$T_\Sigma X \xrightarrow{\text{out}} T(X + \Sigma T_\Sigma X) \xrightarrow{\sigma} S(X + \Sigma T_\Sigma X) \xrightarrow{[\eta \text{inl}, (S \text{inr})v]^*} S(X + T_\Sigma X).$$

In other words, $\mathbb{T}_\Sigma$ is free as an Elgot monad over Σ that extends $\mathbb{T}$.

Example 6.4. Let us spell out what a strong natural transformation $v : \Sigma \rightarrow \mathbb{S}$ amounts to in the case $\Sigma X = \sum_i a_i \times X^{b_i}$ (Example 2.1). A natural transformation $v : \Sigma \rightarrow \mathbb{S}$ is equivalent to a family of natural transformations $a_i \times (-)^{b_i} \rightarrow \mathbb{S}$, equivalently $(-)^{b_i} \rightarrow \mathbb{S}^{a_i}$,

each of which is, by the enriched Yoneda lemma, equivalent to an element of $(Sb_i)^{a_i}$, i.e. a morphism $u_i : a_i \rightarrow Sb_i$. Concretely, v is assembled from the u_i as follows:

$$v_X = [\lambda\langle x, f \rangle. (Sf)(u_1(x)), \dots, \lambda\langle x, f \rangle. (Sf)(u_n(x))].$$

Note that the above generic argument makes use of the assumption that $\mathbf{C}$ is Cartesian closed. In fact it suffices to assume that only the exponentials $(-)^{b_i}$ exist (in particular we do not actually need the exponentials $(-)^{a_i}$ mentioned in between). The expressions $\lambda\langle x, f \rangle. (Sf)(u_i(x))$ above then have to be read as

$$a_i \times X^{b_i} \xrightarrow{\text{swap}} X^{b_i} \times a_i \xrightarrow{\text{id} \times u_i} X^{b_i} \times Sb_i \xrightarrow{\tau} S(X^{b_i} \times b_i) \xrightarrow{S\text{ev}} SX.$$

where swap and ev are the obvious swapping and evaluation transformations respectively.

If $\mathbf{CElg}(\mathbf{C})$ has an initial object, then the statement of Theorem 6.3 can be phrased slightly more concisely. We later give a sufficient criterion on $\mathbf{C}$ that ensures this (Theorem 6.10).

Corollary 6.5. *Suppose that $\mathbf{CElg}(\mathbf{C})$ has an initial object $\mathbb{L}$. Then*

- (1) $\mathbb{L}_\Sigma$ is the free complete Elgot monad over the strong functor $\Sigma : \mathbf{C} \rightarrow \mathbf{C}$, with universal arrow

$$\iota^\mathbb{L} : \Sigma \rightarrow L_\Sigma.$$

- (2) For any complete Elgot monad $\mathbb{T}$, the coinductive generalized resumption monad $\mathbb{T}_\Sigma$ is the coproduct of $\mathbb{T}$ and $\mathbb{L}_\Sigma$ in $\mathbf{CElg}(\mathbf{C})$, with left injection $\text{ext} : \mathbb{T} \rightarrow \mathbb{T}_\Sigma$ and with the right injection being the free extension of $\iota^\mathbb{T} : \Sigma \rightarrow T_\Sigma$ to $\mathbb{L}_\Sigma$.

Proof. Claim (1) is proved by taking $\mathbb{T} = \mathbb{L}$ in Theorem 6.3. Claim (2) is then immediate. $\square$

We assembly some auxiliary results before embarking on the proof of Theorem 6.3.

Lemma 6.6. *The Kleisli composition of a complete Elgot monad $\mathbb{T}$ can be characterized in terms of iteration as follows:*

$$g^* f = [T(\text{inr inr})f, (T \text{ inl}) g]^\dagger \text{ inl} \quad (6.3)$$

Proof. By straightforward calculation:

$$\begin{aligned} & [T(\text{inr inr})f, (T \text{ inl})g]^\dagger \text{ inl} \\ &= [\eta, [T(\text{inr inr})f, (T \text{ inl})g]^\dagger]^* T(\text{inr inr})f && \text{(unfolding)} \\ &= ([\eta, [T(\text{inr inr})f, (T \text{ inl})g]^\dagger] \text{ inr inr})^* f \\ &= ([T(\text{inr inr})f, (T \text{ inl})g]^\dagger \text{ inr})^* f \\ &= ([\eta, [T(\text{inr inr})f, (T \text{ inl})g]^\dagger]^* (T \text{ inl}) g)^* f && \text{(unfolding)} \\ &= g^* f \end{aligned}$$

$\square$

Lemma 6.7. *Let $f : X \rightarrow T(Y + X)$. Then $[\eta, f^\dagger]^* = (T(\text{id} + f))^\dagger$.*

Proof. Consider the following trivially commuting diagram

$$\begin{array}{ccc} X & \xrightarrow{f} & T(Y + X) \\ \downarrow f & & \downarrow T(\text{id} + f) \\ T(Y + X) & \xrightarrow{T(\text{id} + f)} & T(Y + T(Y + X)) \end{array}$$

By uniformity, this implies $f^\dagger = (T(\text{id} + f))^\dagger f$. Therefore $[\eta, f^\dagger]^\star = [\eta, (T(\text{id} + f))^\dagger f]^\star = [\eta, (T(\text{id} + f))^\dagger]^\star T(\text{id} + f) = (T(\text{id} + f))^\dagger$ and we are done. $\square$

We now proceed with the proof of the universal property:

Proof of Theorem 6.3. We first show that ξ has the requisite properties, and then prove uniqueness.

Commutation of Diagram 6.2. We need to show that $\xi \iota^\mathbb{T} = v$ and $\xi \text{ext} = \sigma$. Put $w = [\eta \text{inl}, (S \text{inr})v]^\star \sigma$. Then we have

$$\begin{aligned}
\xi \iota^\mathbb{T} &= \xi \text{out}^{-1} \eta \text{inr} \Sigma \eta^\nu \\
&= (w \text{out})^\dagger \text{out}^{-1} \eta \text{inr} \Sigma \eta^\nu \\
&= [\eta, (w \text{out})^\dagger]^\star w \text{out} \text{out}^{-1} \eta \text{inr} \Sigma \eta^\nu \\
&= [\eta, (w \text{out})^\dagger]^\star [\eta \text{inl}, (S \text{inr})v]^\star \sigma \eta \text{inr} \Sigma \eta^\nu \\
&= [\eta, (w \text{out})^\dagger]^\star [\eta \text{inl}, (S \text{inr})v] \text{inr} \Sigma \eta^\nu \\
&= [\eta, (w \text{out})^\dagger]^\star S(\text{inr})v \Sigma \eta^\nu \\
&= [\eta, \xi]^\star S(\text{inr} \eta^\nu)v \\
&= (\xi \eta^\nu)^\star v \\
&= \eta^\star v \\
&= a.
\end{aligned}$$

and

$$\begin{aligned}
\xi \text{ext} &= ([\eta \text{inl}, (S \text{inr})v]^\star \sigma \text{out})^\dagger \text{ext} \\
&= [\eta, ([\eta \text{inl}, \sigma \text{out}]^\star (S \text{inr})v)^\dagger]^\star \sigma \text{out} \text{ext} && (\text{dinaturality}) \\
&= [\eta, ([\eta \text{inl}, \sigma \text{out}]^\star (S \text{inr})v)^\dagger]^\star \sigma \text{out} \text{out}^{-1} T \text{inl} \\
&= [\eta, ([\eta \text{inl}, \sigma \text{out}]^\star (S \text{inr})v)^\dagger]^\star \sigma T \text{inl} \\
&= [\eta, ([\eta \text{inl}, \sigma \text{out}]^\star (S \text{inr})v)^\dagger]^\star (S \text{inl})\sigma \\
&= \sigma
\end{aligned}$$

ξ is a complete Elgot monad morphism: We have to show that $\xi_X = \zeta_X^\dagger : T_\Sigma X \rightarrow SX$ is natural in X and satisfies the identities (6.1). We successively reduce verification of these properties to the last identity in (6.1), whose proof is the major challenge in establishing the claim.

Note that ζ is a natural transformation (being a composite of natural transformations), and hence $\zeta T_\Sigma f = S(f + T_\Sigma f)\zeta = S(\text{id} + T_\Sigma f)S(f + \text{id})\zeta$ for any f . Therefore, by the uniformity and naturality axioms we obtain

$$\xi T_\Sigma f = \zeta^\dagger T_\Sigma f = (S(f + \text{id})\zeta)^\dagger = (Sf)\zeta^\dagger = (Sf)\xi,$$

i.e. ξ is natural. The equation $\xi\eta = \eta$ is shown as follows:

$$\begin{aligned}\xi\eta &= [\eta, \xi]^*[\eta \text{ inl}, (S \text{ inr})v]^* \sigma \text{ out } \eta \\ &= [\eta, \xi]^*[\eta \text{ inl}, S(\text{inr } f)v]^* \sigma \text{ out } \text{out}^{-1} \eta \text{ inl} \\ &= [\eta, \xi]^* \eta \text{ inl} \\ &= \eta.\end{aligned}$$

Compatibility of ξ with Kleisli lifting follows from Lemma 6.6 and compatibility of ξ with iteration, which we argue later:

$$\begin{aligned}\xi g^* f &= \xi[T_\Sigma(\text{inr inr})f, (T_\Sigma \text{ inl})g]^\dagger \text{ inl} \\ &= [S(\text{inr inr})\xi f, (S \text{ inl})\xi g]^\dagger \text{ inl} \\ &= (\xi g)^*(\xi f).\end{aligned}$$

We now show that $\xi = \zeta^\dagger$ is compatible with strength, i.e. $\xi\tau^\nu = \tau(\text{id} \times \xi)$. With a view to applying uniformity, we calculate $\zeta\tau^\nu$:

$$\begin{aligned}\zeta\tau^\nu &= [\eta \text{ inl}, (S \text{ inr})v]^* \sigma \text{ out } \tau^\nu \\ &= [\eta \text{ inl}, (S \text{ inr})v]^* \sigma T(\text{id} + \Sigma\tau^\nu)(T\delta)\tau(\text{id} \times \text{out}) & (4.3) \\ &= [\eta \text{ inl}, (S \text{ inr})v]^* S(\text{id} + \Sigma\tau^\nu)(S\delta)\tau(\text{id} \times \sigma \text{ out}) & (\sigma \text{ monad morph.}) \\ &= [\eta \text{ inl}, (S \text{ inr})v(\Sigma\tau^\nu)]^* (S\delta)\tau(\text{id} \times \sigma \text{ out}) \\ &= [\eta \text{ inl}, (S \text{ inr})(S\tau^\nu)v]^* (S\delta)\tau(\text{id} \times \sigma \text{ out}) & (\text{naturality of } v) \\ &= [\eta \text{ inl}, S(\text{id} + \tau^\nu)(S \text{ inr})v]^* (S\delta)\tau(\text{id} \times \sigma \text{ out}) \\ &= S(\text{id} + \tau^\nu)[\eta \text{ inl}, (S \text{ inr})v]^* (S\delta)\tau(\text{id} \times \sigma \text{ out}) \\ &= S(\text{id} + \tau^\nu)[\eta \text{ inl}, (S \text{ inr})v]^* S(\text{id} + \rho)(S \text{ dist})\tau(\text{id} \times \sigma \text{ out}) & (\text{def. of } \delta) \\ &= S(\text{id} + \tau^\nu)([\eta \text{ inl}, (S \text{ inr})v\rho] \text{ dist})^* \tau(\text{id} \times \sigma \text{ out}) \\ &= S(\text{id} + \tau^\nu)([\eta \text{ inl}, (S \text{ inr})\tau(\text{id} \times v)] \text{ dist})^* \tau(\text{id} \times \sigma \text{ out}). & (\text{strong nat. of } v)\end{aligned}$$

Furthermore we simplify the tail of the latter expression:

$$\begin{aligned}&([\eta \text{ inl}, (S \text{ inr})\tau(\text{id} \times v)] \text{ dist})^* \tau(\text{id} \times \sigma \text{ out}) \\ &= (S \text{ dist})(S \text{ dist}^{-1})([\eta \text{ inl}, (S \text{ inr})\tau(\text{id} \times v)] \text{ dist})^* \tau(\text{id} \times \sigma \text{ out}) \\ &= (S \text{ dist})([\eta(\text{id} \times \text{inl}), S(\text{id} \times \text{inr})\tau(\text{id} \times v)] \text{ dist})^* \tau(\text{id} \times \sigma \text{ out}) & (\text{def. of } \text{dist}^{-1}) \\ &= (S \text{ dist})([\tau(\text{id} \times \eta \text{ inl}), \tau(\text{id} \times (S \text{ inr})v)] \text{ dist})^* \tau(\text{id} \times \sigma \text{ out}) & (\text{STR}_3) \\ &= (S \text{ dist})(\tau(\text{id} \times [\eta \text{ inl}, (S \text{ inr})v]))^* \tau(\text{id} \times \sigma \text{ out}) \\ &= (S \text{ dist})\tau(\text{id} \times [\eta \text{ inl}, (S \text{ inr})v]^*)(\text{id} \times \sigma \text{ out}) \\ &= (S \text{ dist})\tau(\text{id} \times \zeta).\end{aligned}$$

We have obtained in summary that $\zeta\tau^\nu = S(\text{id} + \tau^\nu)(S \text{ dist})\tau(\text{id} \times \zeta)$. Therefore, by uniformity and compatibility of strength and iteration we obtain the desired identity:

$$\xi\tau^\nu = \zeta^\dagger\tau^\nu = ((S \text{ dist})\tau(\text{id} \times \zeta))^\dagger = \tau(\text{id} \times \zeta^\dagger) = \tau(\text{id} \times \xi).$$

Finally, we are left to show that

$$\xi f^\dagger = (\xi f)^\dagger \tag{6.4}$$

for any $f : X \rightarrow T_\Sigma(Y + X)$ where $\xi = ([\eta \text{ inl}, (S \text{ inr})v]^* \sigma \text{ out})^\dagger$. We proceed by successive reduction of the unrestricted identity (6.4) to the partial cases when f is guarded, and when f is *strongly guarded*. The latter auxiliary notion is defined as follows. Recall that guardedness of f means that $\text{out } f$ factors through some $g : X \rightarrow T(Y + \Sigma T_\Sigma(Y + X))$. We us call f *strongly guarded* if moreover there is $g' : X \rightarrow T(Y + \Sigma X)$ such that $\text{out } f = T(\text{inl} + \Sigma(\eta^\nu \text{ inr}))g'$.

- *Reduction from unrestricted f to guarded f .* Assuming that (6.4) holds for guarded f , we obtain that for any f , $\xi f^\dagger = \xi(\star f)^\dagger = (\xi \star f)^\dagger$. We are left to show that $(\xi \star f)^\dagger = (\xi f)^\dagger$. To that end, consider the morphism w given by the composition

$$X \xrightarrow{\text{out } f} T((Y + X) + \Sigma T_\Sigma(Y + X)) \xrightarrow{[\eta(\text{inl} + \text{id}), (S \text{ inl})\xi^* v]^* \sigma} S((Y + X) + X).$$

Now, on the one hand

$$\begin{aligned} (S[\text{id}, \text{inr}]w)^\dagger &= ([\eta[\text{inl}, \text{inr}], \xi^* v]^* \sigma \text{ out } f)^\dagger \\ &= ([\eta, \xi^* v]^* \sigma \text{ out } f)^\dagger \\ &= ([\eta, \xi]^* [\eta \text{ inl}, (S \text{ inr})v]^* \sigma \text{ out } f)^\dagger \\ &= (\xi f)^\dagger \end{aligned}$$

and on the other hand, by naturality of $_^\dagger$,

$$\begin{aligned} (w^\dagger)^\dagger &= ([\eta(\text{inl} + \text{id}), (S \text{ inl})\xi^* v]^* \sigma \text{ out } f)^\dagger^\dagger \\ &= ([\eta(\text{inl} + \text{id}), (S \text{ inl})\xi^* v, \eta \text{ inr}]^* S[\text{inl} + \text{id}, \text{inl inr}] \sigma \text{ out } f)^\dagger^\dagger \\ &= ([\eta(\text{inl} + \text{id}), (S \text{ inl})\xi^* v, \eta \text{ inr}]^* \sigma (T\pi) \text{ out } f)^\dagger^\dagger \\ &= ([\eta \text{ inl}, \xi^* v]^* (\sigma (T\pi) \text{ out } f)^\dagger)^\dagger && \text{(naturality)} \\ &= ([\eta, \xi^* v]^* S(\text{inl} + \text{id})(\sigma (T\pi) \text{ out } f)^\dagger)^\dagger \\ &= ([\eta, \xi^* v]^* \sigma T(\text{inl} + \text{id})(\sigma (T\pi) \text{ out } f)^\dagger)^\dagger \\ &= ([\eta, \xi^* v]^* \sigma \text{ out } \star f)^\dagger && \text{(definition of } \star) \\ &= ([\eta, \xi]^* [\eta \text{ inl}, (S \text{ inr})v]^* \sigma \text{ out } \star f)^\dagger \\ &= (\xi \star f)^\dagger. && \text{(unfolding for } \xi) \end{aligned}$$

We thus obtain by the codiagonal axiom that

$$(\xi f)^\dagger = (S[\text{id}, \text{inr}]w)^\dagger = (w^\dagger)^\dagger = (\xi \star f)^\dagger.$$

- *Reduction from guarded f to strongly guarded f .* We proceed under the assumption that f is guarded, i.e. $\text{out } f = T(\text{inl} + \text{id})g$ for some $g : X \rightarrow T(Y + \Sigma T_\Sigma(Y + X))$. Let w be the following morphism:

$$\begin{aligned} T_\Sigma(Y + X) &\xrightarrow{\text{out}} T((Y + X) + \Sigma T_\Sigma(Y + X)) \\ &\xrightarrow{[[\eta \text{ inl inl}, [\eta \text{ inl inl}, (S \text{ inl inr})v]^* \sigma g], (S \text{ inr})v]^* \sigma} S((Y + T_\Sigma(Y + X)) + T_\Sigma(Y + X)). \end{aligned}$$

Then, on the one hand, using dinaturality,

$$\begin{aligned}
 (w^\dagger)^\dagger &= \left(([\eta \text{ inl inl}, [\eta \text{ inl inl}, (S \text{ inl inr})v]^* \sigma g], (S \text{ inr})v]^* \sigma \text{ out})^\dagger \right)^\dagger \\
 &= \left([\eta \text{ inl}, [\eta \text{ inl}, (S \text{ inr})v]^* \sigma g]^* ([\eta \text{ inl}, (S \text{ inr})v]^* \sigma \text{ out})^\dagger \right)^\dagger && \text{(naturality)} \\
 &= ([\eta \text{ inl}, [\eta \text{ inl}, (S \text{ inr})v]^* \sigma g]^* \xi)^\dagger && \text{(definition of } \xi) \\
 &= [\eta, ([\eta \text{ inl}, \xi]^* [\eta \text{ inl}, (S \text{ inr})v]^* \sigma g)^\dagger]^* \xi && \text{(dinaturality)} \\
 &= [\eta, ([\eta, \xi]^* [\eta \text{ inl inl}, (S \text{ inr})v]^* \sigma g)^\dagger]^* \xi \\
 &= [\eta, ([\eta, \xi]^* [\eta \text{ inl}, (S \text{ inr})v]^* \sigma \text{ out } f)^\dagger]^* \xi && \text{(definition of } g) \\
 &= [\eta, ([\eta, \xi]^* \xi f)^\dagger]^* \xi && \text{(definition of } \xi) \\
 &= [\eta, (\xi f)^\dagger]^* \xi. && \text{(unfolding)}
 \end{aligned}$$

and hence $(w^\dagger)^\dagger \eta^\nu \text{ inr} = [\eta, (\xi f)^\dagger]^* \xi \eta^\nu \text{ inr} = [\eta, (\xi f)^\dagger]^* \eta^\nu \text{ inr} = (\xi f)^\dagger$. Next we introduce the following morphism t :

$$\begin{aligned}
 T_\Sigma(Y + X) &\xrightarrow{\text{out}} T((Y + X) + \Sigma T_\Sigma(Y + X)) \\
 &\xrightarrow{[[\eta \text{ inl}, g], \eta \text{ inr}]^*} T(Y + \Sigma T_\Sigma(Y + X)) \\
 &\xrightarrow{T(\text{inl} + \Sigma(\eta^\nu \text{ inr}))} T((Y + T_\Sigma(Y + X)) + \Sigma T_\Sigma(Y + T_\Sigma(Y + X))) \\
 &\xrightarrow{\text{out}^{-1}} T_\Sigma(Y + T_\Sigma(Y + X)).
 \end{aligned}$$

By definition, t is strongly guarded, hence $\xi t^\dagger = (\xi t)^\dagger$.

$$\begin{aligned}
 \xi t &= [\eta, \xi]^* \xi t \\
 &= [\eta, \xi]^* [\eta \text{ inl}, (S \text{ inr})v]^* \sigma \text{ out } t \\
 &= [\eta, \xi^* v]^* \sigma \text{ out } t \\
 &= [\eta, \xi^* v]^* S(\text{inl} + \Sigma(\eta^\nu \text{ inr}))[[\eta \text{ inl}, \sigma g], \eta \text{ inr}]^* \sigma \text{ out} \\
 &= [\eta \text{ inl}, \xi^* S(\eta^\nu \text{ inr})v]^* [[\eta \text{ inl}, \sigma g], \eta \text{ inr}]^* \sigma \text{ out} \\
 &= [\eta \text{ inl}, (S \text{ inr})v]^* [[\eta \text{ inl}, \sigma g], \eta \text{ inr}]^* \sigma \text{ out} \\
 &= [[\eta \text{ inl}, [\eta \text{ inl}, (S \text{ inr})v]^* \sigma g], (S \text{ inr})v]^* \sigma \text{ out} \\
 &= S[\text{id}, \text{inr}]w
 \end{aligned}$$

Using the above derived identities and the codiagonal axiom we obtain that

$$(\xi f)^\dagger = (w^\dagger)^\dagger \eta^\nu \text{ inr} = (S[\text{id}, \text{inr}]w)^\dagger \eta^\nu \text{ inr} = (\xi t)^\dagger \eta^\nu \text{ inr} = \xi t^\dagger \eta^\nu \text{ inr}.$$

We are left to show that $\xi t^\dagger \eta^\nu \text{ inr} = \xi f^\dagger$. We strengthen the latter to $t^\dagger = [\eta^\nu, f^\dagger]^*$, which would imply it as follows: $\xi t^\dagger \eta^\nu \text{ inr} = \xi [\eta^\nu, f^\dagger]^* \eta^\nu \text{ inr} = \xi f^\dagger$.

Since t is guarded, we will be done once we show that $[\eta^\nu, f^\dagger]^*$ satisfies the unfolding axiom for $t^\dagger$. It is easy to verify that $\text{out } f^\dagger = T(\text{id} + \Sigma[\eta^\nu, f^\dagger]^*)g$. Then we have

$$\begin{aligned}
 \text{out}[\eta^\nu, f^\dagger]^* &= [\text{out}[\eta^\nu, f^\dagger], \eta \text{ inr } \Sigma[\eta^\nu, f^\dagger]^*]^* \text{out} \\
 &= [[\eta \text{ inl}, \text{out } f^\dagger], \eta \text{ inr } \Sigma[\eta^\nu, f^\dagger]^*]^* \text{out} \\
 &= [[\eta \text{ inl}, T(\text{id} + \Sigma[\eta^\nu, f^\dagger]^*)g], \eta \text{ inr } \Sigma[\eta^\nu, f^\dagger]^*]^* \text{out}
 \end{aligned}$$

while, on the other hand,

$$\begin{aligned}
\text{out } t^\dagger &= \text{out}[\eta^\nu, t^\dagger]^* \text{out}^{-1} T(\text{inl} + \Sigma(\eta^\nu \text{inr})) [[\eta \text{inl}, g], \eta \text{inr}]^* \text{out} \\
&= [\text{out}[\eta^\nu, t^\dagger], \eta \text{inr} \Sigma[\eta^\nu, t^\dagger]^*]^* T(\text{inl} + \Sigma(\eta^\nu \text{inr})) [[\eta \text{inl}, g], \eta \text{inr}]^* \text{out} \\
&= [\text{out } \eta^\nu, \eta \text{inr} \Sigma t^\dagger]^* [[\eta \text{inl}, g], \eta \text{inr}]^* \text{out} \\
&= [[\eta \text{inl}, T(\text{id} + \Sigma t^\dagger)g], \eta \text{inr} \Sigma t^\dagger]^* \text{out}.
\end{aligned}$$

Hence, indeed, $[\eta^\nu, f^\dagger]^* = t^\dagger$.

• *Strongly guarded f .* Finally, let us show (6.4) with strongly guarded f . Suppose that h is such that $\text{out } f = T(\text{inl} + \Sigma(\eta^\nu \text{inr}))h$. Recall that $\xi = ([\eta \text{inl}, (S \text{inr})v]^* \sigma \text{out})^\dagger$. By uniformity, it suffices to show that

$$[\eta \text{inl}, (S \text{inr})v]^* \sigma \text{out } f^\dagger = S(\text{id} + f^\dagger)\xi f.$$

On the one hand,

$$\begin{aligned}
&[\eta \text{inl}, (S \text{inr})v]^* \sigma \text{out } f^\dagger \\
&= [\eta \text{inl}, (S \text{inr})v]^* \sigma \text{out}[\eta^\nu, f^\dagger]^* f \\
&= [\eta \text{inl}, (S \text{inr})v]^* [\text{out}[\eta^\nu, f^\dagger], \eta \text{inr} \Sigma[\eta^\nu, f^\dagger]^*]^* \sigma \text{out } f \\
&= [\eta \text{inl}, (S \text{inr})v]^* [\text{out}[\eta^\nu, f^\dagger], \eta \text{inr} \Sigma[\eta^\nu, f^\dagger]^*]^* S(\text{inl} + \Sigma(\eta^\nu \text{inr}))\sigma h \\
&= [\eta \text{inl}, (S \text{inr})v]^* [\eta \text{inl}, \eta \text{inr} \Sigma f^\dagger]^* \sigma h \\
&= [\eta \text{inl}, S(\text{inr } f^\dagger)v]^* \sigma h.
\end{aligned}$$

and on the other hand,

$$\begin{aligned}
S(\text{id} + f^\dagger)\xi f &= S(\text{id} + f^\dagger)[\eta, \xi]^* [\eta \text{inl}, (S \text{inr})v]^* \sigma \text{out } f \\
&= S(\text{id} + f^\dagger)[\eta, \xi]^* [\eta \text{inl}, (S \text{inr})v]^* \sigma T(\text{inl} + \Sigma(\eta^\nu \text{inr}))h \\
&= S(\text{id} + f^\dagger)[\eta, \xi]^* [\eta \text{inl}, (S \text{inr})v]^* S(\text{inl} + \Sigma(\eta^\nu \text{inr}))\sigma h \\
&= S(\text{id} + f^\dagger)[\eta, \xi]^* [\eta \text{inl} \text{inl}, S(\text{inr } \eta^\nu \text{inr})v]^* \sigma h \\
&= S(\text{id} + f^\dagger)[\eta \text{inl}, \xi^* S(\eta^\nu \text{inr})v]^* \sigma h \\
&= S(\text{id} + f^\dagger)[\eta \text{inl}, (S \text{inr})v]^* \sigma h \\
&= [\eta \text{inl}, S(\text{inr } f^\dagger)v]^* \sigma h.
\end{aligned}$$

This finishes the proof that ξ is a complete Elgot monad morphism.

Uniqueness. Let $\rho : \mathbb{T}_\Sigma \rightarrow \mathbb{S}$ be a complete Elgot monad morphism such that $\sigma = \rho \text{ext}$ and $v = \rho \text{out}^{-1} \eta \text{inr} \Sigma \eta^\nu$. We have to show that $\rho = \xi = ([\eta \text{inl}, (S \text{inr})v]^* \sigma \text{out})^\dagger$. We rewrite the

last term as follows:

$$\begin{aligned}
 & ([\eta \text{ inl}, (S \text{ inr})v]^* \sigma \text{ out})^\dagger \\
 &= ([\eta \text{ inl}, (S \text{ inr})\rho \text{ out}^{-1} \eta \text{ inr} \Sigma \eta^\nu]^* \sigma \text{ out})^\dagger \\
 &= ([\eta \text{ inl}, \rho (T_\Sigma \text{ inr}) \text{ out}^{-1} \eta \text{ inr} \Sigma \eta^\nu]^* \sigma \text{ out})^\dagger \\
 &= ([\eta \text{ inl}, \rho \text{ out}^{-1} T(\text{inr} + (\Sigma T_\Sigma \text{ inr})) \eta \text{ inr} \Sigma \eta^\nu]^* \sigma \text{ out})^\dagger \\
 &= ([\eta \text{ inl}, \rho \text{ out}^{-1} \eta \text{ inr} \Sigma (T_\Sigma \text{ inr}) \Sigma \eta^\nu]^* \sigma \text{ out})^\dagger \\
 &= ([\eta \text{ inl}, \rho \text{ out}^{-1} \eta \text{ inr} \Sigma (\eta^\nu \text{ inr})]^* \sigma \text{ out})^\dagger \\
 &= ((\rho[\eta \text{ inl}, \text{out}^{-1} \eta \text{ inr} \Sigma (\eta^\nu \text{ inr})])^* \rho \text{ ext out})^\dagger \\
 &= (\rho[\eta \text{ inl}, \text{out}^{-1} \eta \text{ inr} \Sigma (\eta^\nu \text{ inr})]^* \text{ext out})^\dagger \\
 &= \rho([\eta \text{ inl}, \text{out}^{-1} \eta \text{ inr} \Sigma (\eta^\nu \text{ inr})]^* \text{ext out})^\dagger.
 \end{aligned}$$

To finish the calculation we have to verify that the term after ρ vanishes. Note that the term under the iteration operator is guarded. Hence, it suffices to show that id satisfies the corresponding characteristic equation for iteration, i.e. that

$$[\eta, \text{id}]^* [\eta \text{ inl}, \text{out}^{-1} \eta \text{ inr} \Sigma (\eta^\nu \text{ inr})]^* \text{ext out} = \text{id}.$$

We reduce the right-hand side to id as follows:

$$\begin{aligned}
 & [\eta, \text{id}]^* [\eta \text{ inl}, \text{out}^{-1} \eta \text{ inr} \Sigma (\eta^\nu \text{ inr})]^* \text{ext out} \\
 &= [\eta, [\eta, \text{id}]^* \text{out}^{-1} \eta \text{ inr} \Sigma (\eta^\nu \text{ inr})]^* \text{ext out} \\
 &= [\eta, \text{out}^{-1} [\text{out}[\eta, \text{id}], \eta \text{ inr} \Sigma [\eta, \text{id}]^*] \eta \text{ inr} \Sigma (\eta^\nu \text{ inr})]^* \text{ext out} \\
 &= [\eta, \text{out}^{-1} \eta \text{ inr}]^* \text{ext out} \\
 &= [\eta, \text{out}^{-1} \eta \text{ inr}]^* \text{out}^{-1} (T \text{ inl}) \text{ out} \\
 &= \text{out}^{-1} [\text{out}[\eta, \text{out}^{-1} \eta \text{ inr}], \eta \text{ inr} \Sigma [\eta, \text{out}^{-1} \eta \text{ inr}]^*] (T \text{ inl}) \text{ out} && (\text{Theorem 4.3}) \\
 &= \text{out}^{-1} (\text{out}[\eta, \text{out}^{-1} \eta \text{ inr}]^* \text{out}) \\
 &= \text{out}^{-1} [\text{out} \eta, \eta \text{ inr}]^* \text{out} \\
 &= \text{out}^{-1} [\eta \text{ inl}, \eta \text{ inr}]^* \text{out} \\
 &= \text{out}^{-1} \text{out} \\
 &= \text{id}.
 \end{aligned}$$

This proves the claim. $\square$

The existence and the exact shape of the initial complete Elgot monad $\mathbb{L}$ mentioned in Corollary 6.5 depend on the properties of the base category. We recall the key definition of a hyperextensive category [2]:

Definition 6.8. A category $\mathbf{C}$ is *hyperextensive* if

- (1) $\mathbf{C}$ has countable coproducts that are *disjoint*, i.e. the pullback of any two distinct injections is an initial object, and *universal*, i.e. stable under pullbacks; and
- (2) in $\mathbf{C}$, subobjects that are coproduct injections are closed under countable disjoint unions; that is, given countably many pairwise disjoint subobjects $A_i \rightarrow B$ that are coproduct injections, their union $\sum_i A_i \rightarrow B$ is again a coproduct injection.

Examples of hyperextensive categories include **Set**, $\omega\mathbf{Cpo}$, and bounded complete metric spaces as well as all presheaf categories [2]. We refer to subobjects whose inclusion morphisms are (binary) coproduct injections as *summands*, and given a summand, we refer to the partner injection of the corresponding binary coproduct as its *coproduct complement* (we will not need uniqueness of complements). In this terminology, summands are closed under pullbacks (i.e. under preimages) and under countable disjoint unions in hyperextensive categories. From countable disjoint unions we obtain unions of chains:

Lemma 6.9. *Let $\mathbf{C}$ be hyperextensive. Then $\mathbf{C}$ has unions of ω -chains of summands; such unions are again summands, and are universal, i.e. stable under pullbacks (and, hence, under products).*

Proof. Any ascending chain of summands can be transformed into a disjoint union of summands: if A_1 and A_2 are summands of X and A_1 is contained in A_2 , then by universality of coproducts, A_1 is also a summand of A_2 so we can replace A_2 with the coproduct complement of A_1 in A_2 , preserving the union. Universality of unions of ascending chains of summands is then inherited from countable disjoint unions. $\square$

Theorem 6.10. *Let $\mathbf{C}$ be hyperextensive and have binary coproducts. Then the monad $\mathbb{L}$ given by $LX = X + 1$ is ω -continuous. Equipped with the arising complete Elgot monad structure according to Theorem 3.7, $\mathbb{L}$ is the initial complete Elgot monad over $\mathbf{C}$.*

(The conditions of the theorem imply our running assumption that $\mathbf{C}$ is distributive [12].)

Remark 6.11. Let us spell out the definition of the iteration operator figuring in the statement of Theorem 6.10 explicitly. Suppose that $e : X \rightarrow L(Y + X)$. Let X_1 be the preimage of Y under e and $e_1 : X_1 \rightarrow Y$ the restriction of e on X_1 ; for every $i \geq 1$ let X_{i+1} be the preimage of X_i under e and let $e_{i+1} : X_{i+1} \rightarrow X_i$ be the restriction of e to X_{i+1} . By universality of finite coproducts, the X_i are pairwise disjoint summands. By stability of summands under countable disjoint unions, the complement of $\sum_i X_i$ in X is a summand which we denote X_∞ . We obtain the presentation $X = \sum_i X_i + X_\infty$. Now $e^\dagger : X \rightarrow LY$ is the universal map induced by the $\eta e_1 \dots e_i : X_i \rightarrow LY$ and $\perp : X_\infty \rightarrow LY$.

Now L clearly admits only very simple recursive definitions: an equation morphism $e : X \rightarrow L(Y + X) = (Y + X) + 1$ essentially defines each variable in X either as a result from Y or as another variable from X or as divergence. In preparation of the proof of Theorem 6.10, the following lemma shows that the solution of all possible such definitions of this shape is, in any complete Elgot monad, uniquely determined by the Elgot monad laws.

Lemma 6.12. *Let $\mathbb{T}$ be a complete Elgot monad over a hyperextensive category $\mathbf{C}$, let $e : X \rightarrow T(Y + X)$, and let $m : Z \rightarrow X$. Then the following holds.*

- (1) *If $em = \eta \text{inl } u$ for some $u : Z \rightarrow Y$ then $e^\dagger m = \eta u$.*
- (2) *If $em = \perp_{Z, Y+X}$ then $e^\dagger m = \perp_{Z, X}$.*
- (3) *If $em = \eta \text{inr } u$ for some $u : Z \rightarrow X$ then $e^\dagger m = e^\dagger u$.*
- (4) *If $em = \eta \text{inr } m u$ for some $u : Z \rightarrow Z$ then $e^\dagger m = \perp_{Z, Y}$.*

That is: If a variable is defined as a result value, then the solution of the recursive definition for that variable is that result value; if a variable is defined as $\perp$, then the solution is $\perp$; if a variable is defined as another variable, then its solution is that of the other variable; and if a set of variables is defined by mutual recursion without any base case and without use of the algebraic operations of the monad, then the solution for all these variables is $\perp$.

Proof. The first three claims are immediate from the unfolding law and coconstancy of $\perp$ (Lemma 3.8). We show the last claim. We have

$$e m = \eta \text{ inr } m u = \eta(\text{id} + m) \text{ inr } u = T(\text{id} + m) \eta \text{ inr } u,$$

which by uniformity implies $e^\dagger m = (\eta \text{ inr } u)^\dagger$. The claim then follows by Lemma 3.9. $\square$

Proof of Theorem 6.10. The base category $\mathbf{C}$ is, a fortiori, extensive. In any extensive category, $\mathbb{L}$ is the partial map classifier for partial morphisms whose domains are summands; we will call such partial morphisms *summand-partial*. Explicitly, a summand-partial morphism f from X to Y is thus a span $X \xleftarrow{m} D \xrightarrow{f} Y$ where m is a summand; the *domain* of f is m or, by abuse of notation, D . By *preimages under f* we mean pullbacks along the map $f : D \rightarrow Y$ in this span.

Thus, the Kleisli category of $\mathbb{L}$ inherits orderings on its hom-sets from the extension ordering on partial functions. The fact that $\mathbf{C}$ has unions of ω -chains of summands which are again summands (Lemma 6.9) then guarantees that these orderings are ω -complete, and since 0 is a summand, they have bottoms $\bullet \leftarrow 0 \rightarrow \bullet$. We have to verify that Kleisli composition for $\mathbb{L}$ is continuous on both sides and that the remaining conditions of Definition 3.4 are satisfied. We will phrase all arguments in terms of summand-partial morphisms.

Continuity of left Kleisli composition: Let g be a summand-partial morphism from Y to Z , and let $(f_i)_{i \in \mathbb{N}}$ be an ascending chain of summand-partial morphisms from X to Y , with domains D_i . Denoting unions and joins of ascending chains by $\sqcup$ and composition of partial morphisms simply by juxtaposition, we have to show that $(\sqcup_i f_i)g = \sqcup_i f_i g$. The only problem here is to show that the domains of the two sides agree. The domain of $f_i g$ is the preimage E_i of D_i under g ; the domain of $\sqcup_i f_i g$ is the union $\sqcup_i E_i$ of the ascending chain $(E_i)_i$; the domain of $\sqcup_i f_i$ is the union $D = \sqcup_i D_i$; and the domain of $(\sqcup_i f_i)g$ is the preimage E of D under g . By universality of unions of ascending chains, $E = \sqcup_i E_i$.

Continuity of right Kleisli composition: Let g be a summand-partial morphism from X to Y with domain C , and let $(f_i)_{i \in \mathbb{N}}$ be an ascending chain of summand-partial morphisms from Y to Z , with domains D_i and supremum f . We have to show $g(\sqcup_i f_i) = \sqcup_i g f_i$; again, we focus only on the domains. The domain of $g f_i$ is the preimage E_i of C under f_i ; the domain of $\sqcup_i g f_i$ is the union $\sqcup_i E_i$; the domain of f is the union $D = \sqcup_i D_i$; and the domain of $g f$ is the preimage E of C under f . By construction, E_i is contained in D_i , and E is contained in D . Moreover, since f_i maps E_i into C , so does f , and hence E_i is also contained in E (by the universal property of E as a pullback). Denoting the restriction of $f : D \rightarrow Y$ to $E \rightarrow C$ by f' , we thus have the diagram

$$\begin{array}{ccccc} E_i & \xrightarrow{\quad} & E & \xrightarrow{f'} & C \\ \downarrow & & \downarrow & & \downarrow \\ D_i & \xrightarrow{\quad} & \sqcup D_i & \xrightarrow{f} & Y \end{array}$$

where the outer rectangle and the right hand square are pullbacks by construction. By the pullback lemma, it follows that the left hand square is also a pullback. By universality of unions of ascending chains of summands, it now follows that $E = \sqcup_i E_i$, as required.

Continuity of the strength: If the Kleisli morphism $f : Y \rightarrow Z + 1$ corresponds to a summand-partial map with domain D , the Kleisli morphism $\tau(\text{id} \times f) : X \times Y \rightarrow Z + 1$

corresponds to a summand-partial map with domain $X \times D$. Continuity of $\tau(\text{id} \times (-))$ is then immediate from stability of unions of ascending chains of summands under products.

Continuity of copairing: Immediate from the fact that generally, $\bigsqcup_i (D_i + E_i) = \bigsqcup_i E_i + \bigsqcup_i D_i$ because unions of ascending chains of summands are defined via coproducts.

Preservation of $\perp$ by left Kleisli composition: The bottom element of the Kleisli hom-set from X to Y is the unique (summand-)partial morphism with domain 0. Left Kleisli composites of this morphism have domains that are pullbacks of 0, which in extensive categories are again 0.

Preservation of $\perp$ by the strength: The domain of the partial morphism corresponding to $\tau(\text{id} \times \perp) : X \rightarrow Y + 1$ is $X \times 0$, which by extensivity (in fact already by distributivity) is 0.

This establishes that $\mathbb{L}$ is ω -continuous, and hence a complete Elgot monad; by the standard construction of least fixpoints in ω -cpo, the iteration operator of $\mathbb{L}$ then has the form described in Remark 6.11. To see initiality of $\mathbb{L}$, let $\mathbb{S}$ be a complete Elgot monad. We need to show existence of a unique complete Elgot monad morphism $\xi : \mathbb{L} \rightarrow \mathbb{S}$. Since ξ must preserve the unit and unproductive divergence $\perp$ (the latter by preservation of iteration), the only candidate is $\xi = [\eta, \perp]$. It remains to show that ξ is a complete Elgot monad morphism. Thanks to the simplicity of the monad structure of $\mathbb{L}$, it is clear that ξ is a strong monad morphism. The main task is to prove preservation of iteration. So let $e : X \rightarrow L(Y + X) = (Y + X) + 1$. We inductively construct infinite sequences $X_1, X_2, \dots$ and $D_1, D_2, \dots$ of summands of X as follows: Like in Remark 6.11, we take X_1 to be the preimage of Y under e , and for $i > 1$ we take X_i to be the preimage of X_{i-1} under e ; similarly, we take D_1 to be the preimage of 1 under e , and for $i > 1$ we take D_i to be the preimage of D_{i-1} under e . By universality of coproducts, the X_i and D_i are pairwise disjoint summands (that is, the X_i are pairwise disjoint, the D_i are pairwise disjoint, and every X_i is disjoint with every D_j).

Let $X' = \sum_i X_i$, $D' = \sum_i D_i$ and let Z be the complement of $X' + D'$ in X . For the remainder we fix the presentation $X = X' + X_\infty$ where $X_\infty = D' + Z$.

By definition, there are $e_1 : X_1 \rightarrow Y$ and $e_i : X_i \rightarrow X_{i-1}$ ($i > 1$) such that

$$e \text{ inl in}_1 = \eta \text{ inl } e_1 \qquad e \text{ inl in}_i = \eta \text{ inr inl in}_{i-1} e_i$$

where in_i denotes the i -th coproduct injection into a countable coproduct.

By applying the unfolding axiom i times we obtain $(\xi e)^\dagger \text{ inl in}_i = \eta e_1 \dots e_i$ and analogously, $\xi e^\dagger \text{ inl in}_i = \xi \eta e_1 \dots e_i = \eta e_1 \dots e_i$. We are left to show that $(\xi e)^\dagger \text{ inr} = \xi e^\dagger \text{ inr}$. Noting that by definition, $e^\dagger \text{ inr} = \perp$ and ξ preserves $\perp$, this amounts to showing that $(\xi e)^\dagger \text{ inr} = \perp$. By construction of the D_i for every $i > 1$ there is $d_i : D_i \rightarrow D_{i-1}$ such that

$$e \text{ inr inl in}_1 = \perp \qquad e \text{ inr inl in}_i = \eta \text{ inr inr inl in}_{i-1} d_i,$$

hence, by applying the unfolding axiom i times we obtain that $(\xi e)^\dagger \text{ inr inl in}_i = \perp$, which implies $(\xi e)^\dagger \text{ inr inl} = \perp$ and hence we are left to show $(\xi e)^\dagger \text{ inr inr} = \perp$.

Notice that the preimages of Y and 1 under $e \text{ inr inr}$ must be 0 and therefore there is $m_1 : Z \rightarrow X$ such that $e \text{ inr inr} = \eta \text{ inr } m_1$. Analogously for every $i > 1$ we construct $m_i : Z \rightarrow X$ such that $e m_{i-1} = \eta \text{ inr } m_i$. Let us denote by $\widehat{Z}$ the sum of ω copies of Z and by $\widehat{m} : \widehat{Z} \rightarrow X$ the cotuple formed by the morphisms m_i with $i > 0$. Now,

$$\begin{aligned} \xi e[\text{inr inr}, \widehat{m}] &= \xi \eta \text{ inr } \widehat{m} w \\ &= \eta \text{ inr}[\text{inr inr}, \widehat{m}] \text{ inr } w \end{aligned}$$

$$= T(\text{id} + [\text{inr inr}, \hat{m}]) \eta \text{inr inr } w$$

where $w : Z + \widehat{Z} \rightarrow \widehat{Z}$ is the obvious canonical isomorphism. By uniformity, and by Lemma 3.9, this implies $(\xi e)^\dagger [\text{inr inr}, \hat{m}] = (\eta \text{inr inr } w)^\dagger = \perp$ and therefore $(\xi e)^\dagger \text{inr inr} = (\xi e)^\dagger [\text{inr inr}, \hat{m}] \text{inl} = \perp$ as required. $\square$

Remark 6.13. The above proof of Theorem 6.10 uses the full power of the definition of hyperextensive categories, including universality of countable coproducts. It has been shown previously [10] that assuming only universality of finite coproducts and stability of summands under countable disjoint unions, one can still define the iteration operator and prove the unfolding axiom. However, we do not see how to show the uniformity axiom in this weaker setting. At the same time, we have the impression that the uniformity axiom is the only place where universality of countable coproducts is needed.

7. EXAMPLE: UNRESTRICTED WHILE LOOPS

We proceed to discuss some examples, aimed mainly at illustrating the benefits of not being restricted to guarded equations in recursive definitions. The first of these is a simple while-language with actions proposed by Rutten, given by the grammar

$$P, Q ::= A \mid P; Q \mid \text{if } b \text{ then } P \text{ else } Q \mid \text{while } b \text{ do } P$$

and, following Piróg and Gibbons [26], interpreted in the Kleisli category of a monad $\mathbb{M}$. Here, A ranges over atomic actions interpreted as Kleisli morphisms $\llbracket A \rrbracket : n \rightarrow Mn$ for some fixed object n , and b over atomic predicates, interpreted as Kleisli morphisms $\llbracket b \rrbracket : n \rightarrow M(1+1)$ (with the left-hand summand read as ‘false’ and the right-hand one as ‘true’). We say that A is of *output type* if $\llbracket A \rrbracket$ has the form $(M \text{fst}) \tau \circ \langle \text{id}_n, p \rangle$ for some $p : n \rightarrow M1$, and of *input type* if $\llbracket A \rrbracket$ factors through $! : n \rightarrow 1$. Sequential composition $P; Q$ is interpreted as Kleisli composition $\llbracket Q \rrbracket^* \llbracket P \rrbracket$, and

$$\llbracket \text{if } b \text{ then } P \text{ else } Q \rrbracket = [\llbracket Q \rrbracket \text{fst}, \llbracket P \rrbracket \text{fst}]^* M \text{dist } \tau \langle \text{id}, \llbracket b \rrbracket \rangle.$$

The key point, of course, is the interpretation of the while loop, given in the presence of iteration $_^\dagger$ by

$$\llbracket \text{while } b \text{ do } P \rrbracket = \left([(M \text{inl}) \circ \eta \text{fst}, (M \text{inr}) \circ \llbracket P \rrbracket \text{fst}]^* M \text{dist } \tau \langle \text{id}, \llbracket b \rrbracket \rangle \right)^\dagger \quad (7.1)$$

where the typing of the expression under the iteration operator $\dagger$ is visualized as

$$\begin{aligned} n &\xrightarrow{\langle \text{id}, \llbracket b \rrbracket \rangle} n \times M(1+1) \\ &\xrightarrow{\tau} M(n \times (1+1)) \\ &\xrightarrow{M \text{dist}} M(n \times 1 + n \times 1) \\ &\xrightarrow{[(M \text{inl}) \circ \eta \text{fst}, (M \text{inr}) \circ \llbracket P \rrbracket \text{fst}]^*} M(n+n). \end{aligned}$$

It has been observed by Piróg and Gibbons that if one instantiates $\mathbb{M}$ with a completely iterative monad, one needs to guard every iteration of the while loop, i.e. change the semantics of while to be

$$\llbracket \text{while } b \text{ do } P \rrbracket = \left([(M \text{inl}) \circ \eta \text{fst}, (M \text{inr}) \circ \llbracket P \rrbracket \text{fst}]^* M \text{dist } \tau \langle \text{id}, \llbracket b \rrbracket \rangle \gamma \right)^\dagger$$

where $\gamma : n \rightarrow Mn$ is guarded, as otherwise the iteration may fail to be defined. If we instantiate $\mathbb{M}$ with an Elgot monad, such as $\mathbb{T}_\Sigma$ for an Elgot monad $\mathbb{T}$, then the guard

is unnecessary, i.e. we can stick to the original semantics (7.1). As an example, consider a simple-minded form of processes that input and output symbols from n and have side effects specified by $\mathbb{T}$; i.e. we work in $\mathbb{M} = \mathbb{T}_\Sigma$ for $\Sigma X = n \times X + X^n$ where we think of Σ as being generated by an output operation $1 \rightarrow n$ and an input operation $n \rightarrow 1$. We correspondingly assume an atomic action *write* that outputs a symbol from n , and an atomic action *read* that inputs a symbol. We interpret *write* as being of output type, i.e. by $\llbracket \text{write} \rrbracket = (M \text{fst}) \tau \circ \langle \text{id}_n, w \rangle$ where $w : n \rightarrow M1$ is obtained from the canonical transformation $\iota^\mathbb{T} : \Sigma \rightarrow T_\Sigma = M$ as defined in Section 6; explicitly, w is the composite

$$n \xrightarrow{\langle \text{id}_n, !_n \rangle} n \times 1 \xrightarrow{\text{inl}} \Sigma 1 \xrightarrow{\iota^\mathbb{T}} M1.$$

Moreover, we interpret *read* as being of input type, i.e. $\llbracket \text{read} \rrbracket = r !_n$ where $r : 1 \rightarrow Mn$ is obtained analogously, i.e. r is the composite

$$1 \xrightarrow{r_0} n^n \xrightarrow{\text{inr}} \Sigma n \xrightarrow{\iota^\mathbb{T}} Mn$$

where $r_0 : 1 \rightarrow n^n$ arises by currying $\text{snd} : 1 \times n \rightarrow n$. Moreover, assume a basic predicate b whose interpretation is largely irrelevant to the example as long as it may take both truth values; for example, b might just pick a truth value non-deterministically or at random, depending on the nature of the base monad $\mathbb{T}$. Consider the program

read; while true do if b then skip else write

where *skip* is an atomic action interpreted as $\llbracket \text{skip} \rrbracket = \eta_n^M : n \rightarrow Mn$. It is possible for the loop to not perform any write operations, as b might happen to always pick the left-hand branch; that is, the loop body fails to be guarded. Since M is an Elgot monad and not just completely iterative, the semantics of the loop is defined (by (7.1)) nonetheless.

8. EXAMPLE: SIMPLE PROCESS ALGEBRA

It has been shown that a simple process algebra BSP featuring finite choice and action prefixing can express all countable transition systems if unguarded recursion is allowed [6, Theorem 5.7.3]. The idea of the proof is to introduce variables X_{ik} for $i, k \in \mathbb{N}$ representing the k -th transition of the i -th state, with X_{i0} representing the i -th state itself, and (unguarded) recursive equations

$$X_{ik} = b_{ik} \cdot X_{j(i,k),0} + X_{i,k+1} \tag{8.1}$$

where the k -th transition of the i -th state performs action b_{ik} and reaches the $j(i,k)$ -th state. (The use of unguarded recursion is essential here, as guarded recursive definitions in BSP will clearly produce only finitely branching systems.) To model this phenomenon using the coinductive generalized resumption transformer, we take $\mathbb{T} = \mathcal{P}_{\omega_1}$, the countable powerset monad on **Set** (see Example 4.6), and the functor Σ generated by a -many unary operations where a is the set of actions; that is, $\Sigma X = a \times X$. We thus regard countable nondeterminism as the base effect, and add action prefixing via coinductive generalized resumptions. Then the definition (8.1) is represented by the map $g = \text{out}^{-1} f : \mathbb{N} \times \mathbb{N} \rightarrow T_\Sigma(\mathbb{N} \times \mathbb{N}) \cong T_\Sigma(0 + \mathbb{N} \times \mathbb{N})$ with $f : \mathbb{N} \times \mathbb{N} \rightarrow T((\mathbb{N} \times \mathbb{N}) + a \times T_\Sigma(\mathbb{N} \times \mathbb{N}))$ given by

$$f(i, k) = \{\text{inr}(b_{ik}, \eta^\nu(j(i, k), 0)), \text{inl}(i, k + 1)\}.$$

Again, our result that $\mathbb{T}_\Sigma$ is an Elgot monad guarantees that this equation has a solution $g^\dagger$; the choice $_^\dagger$ of solutions in $\mathbb{T}_\Sigma$ is uniquely determined as extending the usual structure of $\mathbb{T} = \mathcal{P}_{\omega_1}$ as an Elgot monad, which takes least fixed points.

9. RELATED WORK

The above results benefit from extensive previous work on monad-based axiomatic iteration. In particular we draw on the concept of *Elgot monad* studied by Adámek et al. [3]; the construction of the free Elgot monad over a functor [4] is strongly related to Corollary 6.5.1, and we do not claim this result as a contribution of this paper. There is extensive literature on solutions of (co)recursive program schemes [8, 1, 21, 17, 26, 27], from which our present work differs primarily in that we do not restrict to guarded systems of equations. In particular, as mentioned in the introduction, Piróg and Gibbons [26] actually work with the same monad transformer, the coinductive generalized resumption transformer. Piróg and Gibbons [27, Corollary 4.6] prove a coproduct characterization of the coinductive generalized resumption transformer that is similar to our Theorem 6.3; but again, this takes place in a different category, that is, in completely iterative monads (admitting guarded recursive definitions) rather than complete Elgot monads (admitting unrestricted recursive definitions). Technically, results on T_Σ being a completely iterative monad are incomparable to our result on T_Σ being a complete Elgot monad – we prove a stronger recursion scheme for T_Σ but need to assume that T is an Elgot monad, while T_Σ is completely iterative without any assumptions on T .

We construct solutions of unguarded recursive equations from solutions of guarded recursive equations, for the latter relying crucially on results by Uustalu on guarded recursion over parametrized monads [34], which in particular has allowed us to make do without idealized monads.

The axiomatic treatment of iteration via Elgot monads is essentially dual to the axiomatic treatment of recursion by Simpson and Plotkin [32], who work in a category $\mathbf{D}$ with a *parametrized uniform recursion operator* $\text{Hom}_{\mathbf{D}}(Y \times X, X) \rightarrow \text{Hom}_{\mathbf{D}}(Y, X)$ and a subcategory $\mathbf{S}$ of *strict* functions in $\mathbf{D}$. Given a distributive category $\mathbf{C}$ equipped with a complete Elgot monad, we can take $\mathbf{S} = \mathbf{C}^{op}$ and $\mathbf{D} = (\mathbf{C}_{\mathbb{T}})^{op}$. Then the iteration operator over $\mathbf{C}_{\mathbb{T}}$ sending $f : X \rightarrow T(Y + X)$ to $f^\dagger : X \rightarrow TY$ induces precisely a parametrized uniform recursion operator for the pair $(\mathbf{D}, \mathbf{S})$ in the sense of Simpson and Plotkin.

10. CONCLUSIONS AND FUTURE WORK

We have developed semantic foundations for non-wellfounded side-effecting recursive definitions, specifically for recursive definitions over the so-called *coinductive generalized resumption transformer* that extends a base monad T with operations represented by a functor Σ to obtain a monad T_Σ defined by taking final coalgebras, i.e. consisting of non-wellfounded trees. While previous work on the same monad transformer was focussed on guarded recursive definitions, in the framework of completely iterative monads, we work in the setting of (complete) Elgot monads, which admit unrestricted recursive definitions. Our main results state that

- T_Σ is a complete Elgot monad if T is a complete Elgot monad;
- the structure of T_Σ as a complete Elgot monad is uniquely determined as extending that of T ;
- if the underlying category $\mathbf{C}$ admits an initial complete Elgot monad L (typically $L = _ + 1$) then $T_\Sigma \cong T + L_\Sigma$ in the category of complete Elgot monads on $\mathbf{C}$.

In particular this requires proving the equational laws of complete Elgot monads for the solution operator that we construct on T_Σ . We have implemented a formal verification of our results, which are technically quite involved, in the Coq proof assistant, see <https://git8.cs.fau.de/redmine/projects/corque>.

Besides the fact that applying the coinductive resumption monad transformer to a complete Elgot monad $\mathbb{T}$ again yields a complete Elgot monad $\mathbb{T}_\Sigma$, the resulting object obviously has a richer structure provided by the adjoined free operations. One topic for further investigation is to identify (and possibly axiomatize) this structure. We aim to use this structure to program definitions of free operations as morphisms $T_\Sigma X \rightarrow TX$ in a similar spirit as in the paradigm of *handling algebraic effects* [30]. In conjunction with iteration this actually produces a recursion operator that is more expressive than iteration. This however requires going beyond the first-order setting of this paper (which was sufficient for iteration), as call-by-value recursion is known to be an inherently higher-order concept.

Acknowledgements The authors wish to thank Stefan Milius and Paul Blain Levy for useful discussions.

REFERENCES

- [1] P. Aczel, J. Adámek, S. Milius, and J. Velebil. Infinite trees and completely iterative theories: a coalgebraic view. *Theoret. Comput. Sci.*, 300:1–45, 2003.
- [2] J. Adámek, R. Borger, S. Milius, and J. Velebil. Iterative algebras: How iterative are they? *Theory Appl. Cat.*, 19:61–92, 2008.
- [3] J. Adámek, S. Milius, and J. Velebil. Equational properties of iterative monads. *Inf. Comput.*, 208:1306–1348, 2010.
- [4] J. Adámek, S. Milius, and J. Velebil. Elgot theories: a new perspective of the equational properties of iteration. *Math. Struct. Comput. Sci.*, 21:417–480, 2011.
- [5] J. Adámek and J. Rosický. *Locally Presentable and Accessible Categories*, vol. 189 of *LMS Lect. Notes Ser.* Cambridge Univ. Press, 1994.
- [6] J. Baeten, T. Basten, and M. Reniers. *Process algebra: equational theories of communicating processes*. Cambridge university press, 2010.
- [7] M. Barr. Algebraically compact functors. *J. Pure Appl. Alg.*, 82:211–231, 1992.
- [8] F. Bartels. Generalised coinduction. *Math. Struct. Comput. Sci.*, 13:321–348, 2003.
- [9] J. Bergstra, A. Ponse, and S. Smolka, eds. *Handbook of Process Algebra*. Elsevier, New York, 2001.
- [10] P. Bhaduri and P. Subramanian. Fixed points and iteration in distributive categories. Manuscript, 1997.
- [11] S. Bloom and Z. Ésik. *Iteration Theories*. Springer, 1993.
- [12] A. Carboni, S. Lack, and R. Walters. Introduction to extensive and distributive categories. *J. Pure Appl. Algebra*, 84:145–158, 1993.
- [13] P. Cenciarelli and E. Moggi. A syntactic approach to modularity in denotational semantics. In *Category Theory and Computer Science, CTCS 1993*, 1993.
- [14] R. Cockett. Introduction to distributive categories. *Math. Struct. Comput. Sci.*, 3:277–307, 1993.
- [15] M. Gabbay and A. Pitts. A new approach to abstract syntax involving binders. In *Logic in Computer Science, LICS 1999*, pp. 214–224. IEEE, 1999.
- [16] S. Goncharov, C. Rauch, and L. Schröder. Unguarded recursion on coinductive resumptions. In *Mathematical Foundations of Programming Semantics, MFPS 2015*, ENTCS. Elsevier, 2015.
- [17] S. Goncharov and L. Schröder. A coinductive calculus for asynchronous side-effecting processes. *Inf. Comput.*, 231:204 – 232, 2013.
- [18] M. Hyland, G. Plotkin, and J. Power. Combining effects: Sum and tensor. *Theoret. Comput. Sci.*, 357:70–99, 2006.
- [19] M. Hyland and J. Power. Discrete Lawvere theories and computational effects. *Theoret. Comput. Sci.*, 366:144–162, 2006.
- [20] A. Kock. Strong functors and monoidal monads. *Arch. Math.*, 23:113–120, 1972.

- [21] S. Milius, L. Moss, and D. Schwencke. Abstract GSOS rules and a modular treatment of recursive definitions. *Log. Methods Comput. Sci.*, 9, 2013.
- [22] E. Moggi. An abstract view of programming languages. Technical Report ECS-LFCS-90-113, Univ. of Edinburgh, 1989.
- [23] E. Moggi. A modular approach to denotational semantics. In *Category Theory and Computer Science, CTCS 1991*, vol. 530 of *LNCS*, pp. 138–139. Springer, 1991.
- [24] E. Moggi. Notions of computation and monads. *Inf. Comput.*, 93:55–92, 1991.
- [25] S. Peyton-Jones, ed. *Haskell 98 Language and Libraries — The Revised Report*. Cambridge University Press, 2003. Also: *J. Funct. Prog.* **13** (2003).
- [26] M. Piróg and J. Gibbons. Monads for behaviour. In *Mathematical Foundations of Programming Semantics, MFPS 2013*, vol. 298 of *ENTCS*, pp. 309 – 324, 2013.
- [27] M. Piróg and J. Gibbons. The coinductive resumption monad. In *Mathematical Foundations of Programming Semantics, MFPS 2014*, vol. 308 of *ENTCS*, pp. 273–288, 2014.
- [28] A. Pitts. *Nominal Sets: Names and Symmetry in Computer Science*. Cambridge University Press, 2013.
- [29] G. Plotkin and J. Power. Notions of computation determine monads. In *Foundations of Software Science and Computation Structures, FoSSaCS 2002*, vol. 2303 of *LNCS*, pp. 342–356. Springer, 2002.
- [30] G. Plotkin and M. Pretnar. Handling algebraic effects. *Log. Meth. Comput. Sci.*, 9, 2013.
- [31] J. Rutten. A note on coinduction and weak bisimilarity for while programs. *ITA*, 33:393–400, 1999.
- [32] A. Simpson and G. Plotkin. Complete axioms for categorical fixed-point operators. In *Logic in Computer Science, LICS 2000*, pp. 30–41. IEEE, 2000.
- [33] M. Smyth and G. Plotkin. The category-theoretic solution of recursive domain equations. In *Foundations of Computer Science, FOCS 1977*, pp. 13–17. IEEE Computer Society, 1977.
- [34] T. Uustalu. Generalizing substitution. *ITA*, 37(4):315–336, 2003.
- [35] P. Wadler. How to declare an imperative. *ACM Comput. Surv.*, 29:240–263, 1997.
- [36] M. Wand. Fixed-point constructions in order-enriched categories. *Theoret. Comput. Sci.*, 8:13–30, 1979.